

GROUND-WATER RESOURCES OF THE BALMORHEA AREA
IN WESTERN TEXAS

by

Walter N. White - Hoyt S. Gale and S. Spencer Nye

Report prepared in cooperation between the
Geological Survey, United States Department
of the Interior and the Texas Board of
Water Engineers.

February 1938

CONTENTS

	Page
Introduction.	1
Purpose and scope of the investigation	1
Acknowledgments.	3
Topography and drainage	3
Springs.	4
Climate.	5
General geology	10
The relation of the Balmorhea springs to the geology.	16
Occurrence of ground water in the Balmorhea area.	22
Springs.	22
Artesian springs.	22
Gravity springs	28
Shallow ground water.	31
Records of wells	32
Measurements of water levels in wells.	34
Intake of ground water.	41
Areas of intake.	41
Intake from streams.	43
Madera Canyon.	44
Cherry Canyon.	46
Big Aguja Canyon	46
Little Aguja Canyon.	47
Limpia Creek.	48
Discharge of ground water.	49
Spring discharge.	49
Water analyses.	27-50a
Conclusions.	51

Tables

Analyses of water from large springs	27
Partial analyses of water from gravity springs	50a
Partial analyses of water from wells	50a
Miscellaneous measurements of spring discharge	55
Discharge of Phantom Lake Spring	61-63
Discharge of San Solomon Spring.	65-67
Discharge of Giffin Spring	63-64a
Discharge of West Sandia.	72-74
Discharge measurements in Madera Canyon.	56-57a
Discharge measurements in Cherry Canyon.	74a
Discharge measurements in Big Aguja Canyon	60
Discharge measurements in Little Aguja Canyon.	59
Discharge measurements of Toyah Creek.	58
Discharge measurements of Limpia Creek	75-76
Discharge measurements of canals and laterals of Reeves	
County Water Improvement District No. 1	68-71
Miscellaneous discharge measurements	78-80

ILLUSTRATIONS

	Page
Plate 1. Geologic map and section of Balmorhea area, Texas, showing the relations between the large springs and geologic structure	81
Figure 1. Map of Toyah Basin, Texas, showing springs and areas of ground-water intake near Balmorhea	82
Figure 2. Map of the vicinity of Balmorhea, Texas, showing wells and springs and approximate shape and slope of water table	83

ABSTRACT

Balmorhea, located in the Toyah Basin at the foot of the Davis Mountains in western Texas, is the center of a thriving farming community, the lands of which are irrigated with water derived for the most part from large springs. During prosperous farming years the total area under irrigation has reached about 10,000 acres. In 1932, a year of financial depression and low prices for farm products, the area irrigated was estimated as about 7,000 acres. The irrigation system includes a reservoir which is used to store flood waters from Toyah Creek and a part of the winter flow of the springs. The capacity of the reservoir, originally about 6,000 acre-feet, has been reduced to some extent by silting and partly on this account a shortage of water occurs during the late summer and fall in years when the requirements for irrigation are large. This condition would be relieved and possibly new land could be irrigated if a large additional supply of water could be obtained from wells. On the other hand, the community would suffer severe loss if withdrawals of water through wells should result in a material reduction in the discharge of the springs. One of the chief purposes of the investigation discussed in this report was to obtain information that will throw light on this important question.

The rocks exposed in the vicinity of Balmorhea comprise Lower and Upper Cretaceous marine sediments, Tertiary volcanic deposits and lava flows, and Quaternary alluvial deposits. The Lower Cretaceous rocks consist mainly of massive thick-bedded limestone with some inter-bedded calcareous shale and a basal unit composed of sand or sandstone. Their total thickness at Balmorhea is not known but is believed to be about 500 feet. The Upper Cretaceous rocks are mostly of clayey composition but include some thin bedded flaggy or hard platy limestone at the base. The evidence suggests that these rocks have a thickness of at least 500 or 600 feet in most places. Volcanic rocks of Tertiary age form the capping of the Davis Mountains and of numerous ridges and hills on their flanks in the vicinity of Balmorhea.

Usually the base of the volcanic series consists of a bed of white tuff or ash which is overlain in turn by volcanic breccia and a succession of lava flows with interbedded tuff. These rocks have a total thickness of 1,500 to 2,000 feet. Deposits of gravel and alluvium of late Tertiary, Pliocene and Recent age overlie the bed rock of the lower mountain slopes and most of the lowlands in this vicinity. Above the settlement of Brogada these gravels, according to well records, are 50 feet deep or slightly deeper in a few places but usually are from 15 to 25 feet deep. Below Brogada they are much deeper.

The climate of Balmorhea and the surrounding region is semi-arid and in most years farming can not be carried on successfully without irrigation.

The Balmorhea springs consist of six springs or groups of springs that appear at irregular intervals in the floor of the valley of Toyah Creek, called Phantom Lake, San Solomon, Giffin, Saragosa, West Sandia and East Sandia springs, and a series of springs and seeps that rise in the bed of Toyah Creek. The springs are of two types, artesian and gravity. The largest, San Solomon, Phantom Lake and Giffin which together contribute most of the available water supply belong to the first class and the others to the latter class. It is believed that the artesian springs come from a reservoir in an extensive interconnected system of fissures and solution passages in Lower Cretaceous limestone. The water enters the limestone between the springs and the Davis Mountains, the principal area of intake being a long anticlinal valley which parallels the steep front of the mountains west, northwest and southwest of the springs. In this valley the beveled edges of the limestone appear at the surface or lie beneath a relatively shallow mantle of stream and terrace gravel. The streams from Aguja, Little Aguja, Madera and Cherry Canyon lose heavily between the mouths of their canyons and the down stream edge of this valley. Rainfall on the mountain slopes and bordering gravel covered terraces and seeps that issue at the base of the volcanic rocks also contribute, the water from these sources reaching the limestones by lateral movement through the gravels.

The water rises in the springs by artesian pressure along faults in which relatively impermeable Upper Cretaceous rocks have been dropped down against the water-bearing Lower Cretaceous limestone.

The gravity springs are fed from shallow ground waters in the Toyah Valley gravels. These waters are replenished by rainfall on the valley floor and by seepage from Toyah Creek, canals, ditches and irrigated lands and to some extent by seepage from the artesian springs.

Between the fall of 1931 and the fall of 1933 continuous records of the discharge of San Solomon, Phantom Lake, Giffin and East and West Sandia springs were obtained, and the discharge of Saragosa and Toyah Creek springs was measured occasionally. During the period, the combined flow of the five springs first mentioned was at the rate of about 48,000 acre-feet a year and the combined discharge of the Saragosa and Toyah Creek springs apparently was at the rate of 6,500 to 7,000 acre-feet a year. Thus the estimated flow of all the springs from the fall of 1931 to the fall of 1933 was at the average rate of around 55,000 acre-feet a year. The total rainfall during the two years was considerably above normal, and on that account, the annual discharge is believed to have been materially greater than the long-term average.

Following heavy rains in September 1932, the discharge of San Solomon Spring was more than doubled, and that of Phantom Lake Spring was multiplied about nine times. Also the discharge of Giffin Spring was increased about $33\frac{1}{3}$ per cent, Saragosa Spring about 100 per cent, and Toyah Creek springs about 400 per cent. The flow of East Sandia and West Sandia, however, increased very little. San Solomon and Giffin springs showed an increase almost immediately after the start of the heavy rains. The increase in the flow of these two springs was accompanied by a sharp decline in the temperature and mineral content of the water, and the discharge of both springs became muddy or cloudy for a time.

It is concluded that a well put down in the vicinity of Phantom Lake Spring or in the area between that spring and the San Solomon-Giffin Spring fault might tap the limestone reservoir at shallow depths, but if pumped heavily, it would be practically certain to deplete the flow of the artesian springs. From the San Solomon-Giffin Spring fault to the Brogada Hills, the water-bearing horizon in the Lower Cretaceous limestone probably is around 700 to 800 feet below the surface. If the fault barrier is not complete, a well in that area might deplete the flow of the springs. If the barrier is complete, solution channeling may not have developed extensively in the limestone, because of lack of circulation, and as a result the limestone may be comparatively tight and the water in it highly mineralized. In the district between the Brogada Hills and Saragosa the limestone is perhaps 1,000 to 1,200 feet below the surface, and in that district it is reasonably probable that any water which occurs in the limestone is shut off from the reservoir that supplies the artesian springs. A deep well in that district therefore would not be expected to interfere with their flow.

Some water probably could be developed from shallow wells if they were put down near the upper end of the Balmorhea district, in areas fairly remote from the gravity springs, (Toyah Creek, Saragosa and Sandia), so as to avoid depleting their flow, at least during the irrigation season.

Introduction

Purpose and scope of the investigation

Balmorhea is the center of a thriving farming community the lands of which are irrigated with water derived for the most part from large springs. During prosperous years the total area under irrigation has been about 10,000 acres. In 1932, a year of depression and low prices, the irrigated area was estimated at about 7,000 acres. The Reeves County Improvement District No. 1, a cooperative organization, embracing most of the farms in the area, has a reservoir (see Fig. 2) which is used to store a part of the winter flow of the springs, together with flood waters of Toyah Creek when such waters are available. The reservoir originally had a capacity of about 6,000 acre-feet but silting has reduced the capacity of the reservoir so that a considerable part of the winter flow of the springs and flood water passes the reservoir. As a result there is a shortage of water during the late summer and fall in years when the irrigated acreage is large. The shortage is not serious enough to cause destruction of crops but is sufficient to reduce the yield from some of the lands.

The community would be materially benefited if water could be withdrawn by wells from the natural underground reservoirs without depleting the flow of the springs. Thereby the shortage during the latter part of the irrigation season would be relieved and additional lands might be irrigated. On the other hand, the community might suffer serious loss if the operation of wells should materially reduce the flow of the springs. The loss might be irreparable if wells were put down by outside interests for the irrigation of new lands and the owners of the springs had no recourse.

The chief purpose of the investigation in the Balmorhea district, therefore, was to obtain information that will throw light on this important problem. An important part of the investigation was to compile an authoritative long-time record of the flow of the springs which will serve as evidence, if necessary, to protect the rights of the owners of the springs.

The field work, which was done chiefly in 1931 to 1933, included the following: The geology of the Davis and Barrilla Mountains and Balmorhea district was studied. Weirs and automatic water-stage recorders were maintained for about two years on Phantom Lake, San Solomon, Giffin, West Sandia and East Sandia springs, and continuous records were thereby obtained. Gages were installed on Limpia Creek and the streams in Cherry, Madera, Big Aguja and Little Aguja Canyon, and a water-stage recorder was maintained in Madera Canyon. Several series of measurements were made with current meters along these streams in order to determine losses by seepage and gains by ground-water inflow in various relatively short stretches. Discharge measurements were made to determine seepage losses from the main canal and laterals of the Reeves County Water Improvement District No. 1. The water table in the valley of Toyah Creek near Balmorhea was mapped. The depths to the water levels in 13 selected shallow wells were measured monthly and sometimes weekly for nearly two years and a water-stage recorder was maintained on one of the wells for about 1-1/2 years. Analyses were made of the waters of the springs and selected wells.

In general the geological parts of this report were prepared by Hoyt S. Gale and the discussion of the ground-water conditions by Walter N. White. The preliminary studies of the geology and ground-water conditions of the district and surrounding region were made by S. S. Nye. The measurements of the flow of the springs and streams were largely made by V. W. Rupp.

The investigation was made under the general direction of O. F. Meinzer, geologist in charge of the ground water division of the Geological Survey.

Acknowledgments

The writers are indebted to H. P. Bybee, of the University of Texas, and to P. B. King, of the Geological Survey of the United States Department of the Interior, for geological information and criticism, and to J. J. Bayles, in charge of the Texas Agricultural Farm near Balmorhea, who furnished climatological data and other helpful information. Special acknowledgments are due the Board of Directors of the Reeves County Water Improvement District No. 1 for their cooperation. Mr. A. C. Shreyer gave valuable assistance and advice. Mr. Edgar George and Mr. T. M. Delaney gave efficient assistance in the field work and otherwise expedited the investigation.

Topography and drainage

Balmorhea has an altitude of about 3,200 feet. It is situated along the southwestern margin of the Toyah Basin outwash plain, on Toyah Creek, near the foot of the Davis and Barrilla Mountains (see fig. 1). These mountains are characterized by flat tops, steep slopes and narrow V-shaped canyons. The tops of the Davis Mountains are mostly from 5,000 to 6,000 feet above sea level but Star Mountain and Timber Mountain, two of the most conspicuous flat-topped mountains in this part of the range, rise to 6,350 and 6,442 feet respectively. The highest peak in the Barrilla Range reaches an altitude of 5,560 feet. The surface between the mountains and Balmorhea and around Balmorhea is broken by several low ridges or isolated hills. These include a ridge or line of low hills between Balmorhea and the Davis Mountains which is parallel to the front of the mountains and a narrow northwest-southeast ridge about three miles long east of Balmorhea which is known as the Brogada Hills. The mountains and adjacent plain are drained by Toyah Creek, which is formed about six miles southwest of Balmorhea

by the junction of streams from Big Aguja, Little Aguja and Madera Canyons. Toyah Creek carries great quantities of storm water at times but is generally dry, except a short stretch of about four miles near Balmorhea, where there is a perennial flow of spring water. Most of the storm water is diverted, near Balmorhea, into a reservoir belonging to Reeves County Water Improvement District No. 1. During exceptionally heavy floods a part of the water passes the reservoir diversion dam and either seeps into the gravels of the outwash plain or empties into Toyah Lake, located about 30 miles to the north. The adjacent area on the north is drained by a creek from Cherry Canyon and that on the south is drained by Limpia Creek (sometimes called Barrilla Draw). Both have perennial flow in the mountains but lose their water soon after reaching the foot hills. Both carry large amounts of flood water at times.

Springs

The Toyah Basin is noted for two groups of large springs, one of which is located near Balmorhea and the other near Fort Stockton. The Balmorhea group consists of six springs or groups of springs that appear at irregular intervals in the floor of the valley of Toyah Creek, called Phantom Lake, and San Solomon, Giffin, Saragosa, West Sandia, and East Sandia Springs, and a group of springs and seeps that rise in the bed of Toyah Creek. The location of the springs is shown on plate 1 and figure 1 and 2.

Phantom Lake spring flows from a crevice in limestone at the base of a ridge about eight miles southwest of Balmorhea. San Solomon and Giffin springs rise from gravel deposits in the floor of the valley near Toyahvale, about four miles southwest of Balmorhea. Saragosa, West Sandia and East Sandia springs rise from gravel in the bottom of shallow drainage channels near Balmorhea. The Toyah Creek springs rise in gravels in the

channel of the creek near Balmorhea.

The springs have a combined discharge of about 23,000 gallons a minute during dry years and a much larger discharge during wet years. Measurements of the discharge of all the springs are given on pp 55-80. San Solomon Spring, Phantom Lake and Giffin Springs rise from a vast reservoir in cavernous limestone. The others have their source in relatively shallow sand and gravel and are essentially water-table springs.

Climate

The climate of Balmorhea and the adjacent region is semi-arid, and in most years farming cannot be carried on successfully without irrigation. In summer the days are hot, but the nights are cool. The winters are comparatively mild, but killing frosts are not uncommon between November and March.

The U. S. Weather Bureau has obtained records of precipitation at the State Agricultural Experiment Station, near Balmorhea, at Fort Davis, 30 miles south of Balmorhea, during 55 years since 1857, and at Fort Stockton, 50 miles east of Balmorhea, during 60 years since 1870. The average annual precipitation during the periods of observation down to 1936, inclusive, was as follows: Balmorhea, 13.43 inches; Fort Davis, 16.51 inches; and Fort Stockton, 14.89 inches.

In most years between 1924 and 1936 the precipitation at the three stations showed similar trends, but there were exceptions to that general rule. For example, in 1933 the precipitation was about the average at Fort Stockton and much below the average at Balmorhea and Fort Davis, and in 1936 it was well above the average at Fort Davis, about the average at Balmorhea, and below the average at Fort Stockton. If the long-time records at Fort Davis and Fort Stockton are accepted as a basis for estimate, it would appear that the average annual precipitation at Balmorhea from 1924

to 1936 did not differ materially from the long time average in that vicinity, but that the proportion of very dry years, namely four out of 13, was unusually high. Exceptional years in the Balmorhea record were 1932, with a precipitation of 28.15 inches, 1933, with a precipitation of only 6.43 inches, and 1934, with a precipitation of 3.89 inches.

The precipitation in the Davis Mountains averages considerably higher than the average at Balmorhea, but no accurate information is available from which a comparison can be made. Some pine timber is found on the tops of the mountains, notably on Timber Mountain, but the mountain vegetation generally consists of cedars and brush and shrubs that are common to semi-arid regions. According to common report the average rainfall is higher in the mountains, but the storms, particularly in summer, are erratic. It often rains in the mountains when none falls at Balmorhea and sometimes the opposite occurs. Rains occur at times on a part of the mountains and produce a run of storm water in one canyon while on adjacent areas the canyon streams remain dry.

The following tables, taken from the records of the United States Weather Bureau, give the annual precipitation at Fort Davis from 1855 to 1860, from 1870 to 1890 and from 1902 to 1936; at Fort Stockton from 1870 to 1886 and from 1894 to 1936; the monthly precipitation at Balmorhea from October 1923 to December 1936, and the daily precipitation at Balmorhea from October 1931 to September 30, 1933.

Annual precipitation in inches
at Fort Davis, Jeff Davis County, Texas

Year	Precipitation	Year	Precipitation	Year	Precipitation
1855	21.21	1883	14.22	1919	20.22
1856	25.94	1884	22.56	1920	25.11
1857	19.82	1885	14.22	1921	10.10
1858	14.12	1886	12.64	1922	12.33
1859	22.55	1887	18.50	1923	13.17
1860	8.52	1888	18.11	1924	14.14
1870	12.67	1889	11.34	1925	16.65
1871	6.78	1890	18.34	1927	13.09
1872	10.01	1902	14.95	1928	14.30
1873	16.29	1903	12.39	1929	14.03
1874	20.10	1904	20.19	1930	15.80
1875	27.68	1905	23.13	1931	15.54
1876	23.92	1912	12.93	1932	25.56
1877	16.16	1913	18.82	1933	11.73
1878	15.43	1914	23.95	1934	9.40
1879	21.41	1915	11.98	1935	10.87
1880	23.48	1916	10.10	1936	20.37
1881	27.54	1917	9.35		
1882	20.22	1918	14.30	Average	16.51

Annual precipitation in inches
At Fort Stockton, Pecos County, Texas

Year	Precipitation	Year	Precipitation	Year	Precipitation
1870	18.58	1897	11.21	1917	5.68
1871	5.88	1898	7.29	1918	14.54
1872	12.99	1899	10.22	1919	24.79
1873	11.20	1900	19.58	1920	21.39
1874	13.90	1901	10.17	1921	10.32
1875	16.78	1902	11.42	1922	12.11
1876	12.45	1903	12.00	1923	18.83
1877	13.00	1904	15.97	1924	9.82
1878	12.47	1905	20.65	1925	20.43
1879	5.12	1906	19.27	1926	18.32
1880	33.76	1907	12.01	1927	12.12
1881	12.65	1908	12.31	1928	18.30
1882	25.56	1909	11.27	1929	16.30
1883	27.39	1910	4.07	1930	9.00
1884	24.07	1911	20.24	1931	15.97
1885	20.02	1912	10.31	1932	24.61
1886	9.84	1913	13.11	1933	14.63
1894	10.73	1914	22.58	1934	6.87
1895	27.70	1915	13.92	1935	10.12
1896	16.33	1916	7.95	1936	12.75
				Average	14.89

Monthly precipitation in inches at Balmorhea, Reeves County, Texas, 1923-1936.

Year	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.	Oct.	Nov.	Dec.	Annual
1923									.90	1.48	1.48	2.17	
1924	T.	1.98	.33	.46	.02	T.	1.50	.21	2.64	1.74	T.	.23	9.11
1925	T.	.00	.00	2.22	1.43	.13	2.13	3.73	2.27	2.65	T.	.09	14.65
1926	.60	.00	1.68	.75	1.50	1.36	2.60	2.68	2.09	2.69	.74	1.62	18.31
1927	.13	.44	.63	T.	.37	.54	.94	.46	3.74	.37	.00	.58	8.20
1928	.01	.12	.03	.39	1.33	.66	4.97	4.62	3.06	.93	.64	.12	16.88
1929	.00	.77	.03	.84	1.28	1.33	.59	.92	2.13	1.81	.33	.23	11.16
1930	.32	.16	.47	1.50	.93	4.87	.72	2.82	.47	1.98	.85	.73	15.82
1931	1.13	.90	.89	3.47	1.35	1.69	1.25	1.47	.43	.21	.86	2.25	15.90
1932	.48	3.79	.58	.45	2.19	.48	1.55	3.60	11.64	1.20	T.	2.19	28.15
1933	.08	.61	T.	.03	.22	.42	.60	1.09	.92	2.20	.23	.03	6.43
1934	.39	.07	.29	.32	.23	1.03	.52	.81	.14	T.	.08	.01	3.89
1935	T.	.71	T.	.45	1.58	.97	.50	3.20	2.57	1.06	.34	1.95	12.45
1936	.54	.00	.82	.09	2.65	.48	.95	.59	5.15	.56	1.81	.11	13.75
Average	.28	.73	.55	.85	1.16	1.07	1.45	2.01	2.72	1.34	.52	.87	13.43

Rainfall recorded at Balmorhea, Texas

October 1, 1931 to September 30, 1933, inclusive
(inches)

1931		1932, Cont'd.		1932, Cont'd.		1933, Cont'd.	
Oct. 11	.06	Apr. 26	.27	Sept. 7	1.83	Mar.	Trace
12	.09	27	.18	8	.44		
13	.06	Total	0.45	19	.01	Apr. 25	.03
Total	0.23			22	.26	Total	.03
		May 10	.33	23	.16		
Nov. 16	.03	11	.34	24	.03	May 17	.08
17	.21	14	.14	25	.13	24	.04
25	.01	16	.03	26	.44	25	.10
26	.05	24	.48	27	.02	Total	0.22
30	.56	25	.35	28	.35		
Total	0.86	27	.10	29	3.30	June 2	.13
		29	.40	30	.47	13	.05
Dec. 1	.92	Total	2.19	Total	11.64	19	.05
2	.15					29	.19
7	.47	June 19	.20	Oct. 1	.07	Total	0.42
8	.12	22	.28	2	.07		
17	.15	Total	0.48	10	.05	July 17	.09
18	.44			13	.39	18	.27
Total	2.25	July 3	.17	14	.19	19	.18
		4	.14	15	.01	31	.06
		7	.70	20	.09	Total	0.60
		8	.03	21	.28		
1932		9	.03	25	.05	Aug. 2	.02
Jan. 4	.03	10	.07	Total	1.20	12	.09
11	.34	22	.26			16	.04
15	.08	26	.05	Nov. Trace		24	.08
22	.03	30	.05			26	.04
Total	0.48	Total	1.55	Dec. 9	.13	27	.06
				10	.05	28	.04
Feb. 14	.13	Aug. 7	.21	14	.34	29	.58
15	.37	8	.04	15	.13	30	.13
17	.03	9	.12	17	.07	31	.01
18	.17	12	.20	22	.07	Total	1.09
19	.40	18	.16	23	.91		
20	.75	26	.07	29	.05	Sept. 1	.75
21	1.06	27	.03	30	.46	2	.01
22	.80	29	.05	Total	2.19	9	.05
25	.08	30	2.72			13	.04
Total	3.79	Total	3.60			14	.07
				1933		Total	0.92
Mar. 2	.01	Sept. 2	1.03	Jan. 7	.06		
4	.38	3	.05	8	.02		
5	.14	4	2.63	Total	0.08		
12	.05	5	.10				
Total	0.58	6	.39	Feb. 8	.05		
				25	.28		
				26	.28		
				Total	0.61		

General geology
By Hoyt S. Gale

The rocks exposed in the vicinity of Balmorhea comprise Lower and Upper Cretaceous marine sediments, Tertiary volcanic deposits and lava flows, and Quaternary alluvial deposits. The Cretaceous sediments are probably underlain by a succession of Triassic, Permian and older strata that are exposed in adjoining areas, particularly in the Delaware and Guadalupe Mountains to the northwest and in the Glass Mountains and Marathon area to the southeast. The occurrence of these rocks has been extensively studied in recent years by geologists working for the various oil companies, and a large amount of information has been published concerning them. A review of this material, and field study centering in the Toyah Basin, including the Balmorhea area, by the writer in the spring and fall of 1932 and a part of the spring of 1933 is the basis for the following summary statement of the geology of the area.

Lower Cretaceous rocks.- It is believed that the big springs of Balmorhea and Fort Stockton come from an extensive network of fissures and solution passages in limestones of the Comanchean series of Lower Cretaceous age, which are the oldest strata exposed near Balmorhea. The Comanche section includes, besides the limestones, interbedded calcareous shale and a basal unit composed of sand or sandstone, at places containing gravel, which although not exposed near Balmorhea is found in outcrops near Kent to the northwest and in the foothills of the Glass Mountains near Hovey to the southeast. The limestones are mainly massive, thick-bedded and very fossiliferous. They characteristically weather to dark-colored craggy or solution-rounded ledges, and low rolling to hilly topography.

The total thickness of the Lower Cretaceous at Balmorhea is not known because the base of the series is not exposed near that place and as far as

is known none of the wells that have been drilled in the area have penetrated the series. Near Fort Stockton, east of Balmorhea, it has been shown that the maximum thickness of the series is about 600 feet, and near Limpia post-office, 16 miles due south of Balmorhea, a deep well passed through about 700 feet of Lower Cretaceous rocks. It is believed that the Lower Cretaceous section thins toward the northwest, and that in the vicinity of Balmorhea the series is approximately 500 feet thick.

The uppermost limestone of the Lower Cretaceous series is well exposed near Phantom Spring in a ridge that extends northwest from the spring along the south side of the Old Spanish Trail highway. This limestone contains several "sinks" or deep cavernous channels and crevices in this general vicinity and from one of these crevices emerges the water that feeds Phantom Lake. The section is repeated in a parallel ridge on the north side of the highway by a fault that passes near the spring (see map, plate 1, p. 81). This area of outcrop of the Lower Cretaceous extends essentially continuously from these exposures to and beyond the Texas and Pacific Railroad at San Martine in a gradually broadening band which is structurally a faulted anticline that plunges toward the southeast. A second faulted anticline, parallel to the front of the Davis Mountains and to the first-mentioned anticline, passes through the summit of Star Mountain and brings the limestone to the surface again in a large area in the hills at the foot of the steep slope east of Gomez Peak (fig. 1, p. 82). This anticline has been carved by erosion into a broad valley which is adjacent to the steep front of the mountains and lies athwart the courses of all the streams from Cherry Creek to Aguja Creek. The anticline plunges to the southeast and the Lower Cretaceous limestone finally disappears beneath the stream deposits that underlie the surface in most places in the valley area. The southeasternmost exposure of the Lower

Cretaceous along this anticline is found in a small outcrop in the channel of Madera Creek five or six miles southwest of Phantom Lake Spring where the creek crosses the axis of the anticline. Cherry, Madera, Aguja and Little Aguja Creeks and other smaller streams that cross the anticline lose heavily or disappear altogether into the gravel. It is believed that this water passes through the gravel and enters the Lower Cretaceous rocks that doubtless closely underlie the gravel a part of the way across the anticlinal valley.

There is a very extensive exposure of Lower Cretaceous limestone southeast of Balmorhea and east and southeast of the Barrilla Mountains in an area that is crossed by Limpia Creek and other streams from the north slope of the Davis Mountains. Limpia Creek disappears entirely during moderate or low stages in that part of its course and in a stretch underlain by gravels immediately above the outcrop. The Lower Cretaceous is also exposed in an area in the center of the main anticline of the Barrilla Mountains southeast of Balmorhea and may take in considerable water there.

Upper Cretaceous rocks. - The Upper Cretaceous rocks in this area are mostly of clayey composition, but they include some limestones in the lower part of the series that may be subject to solution channelling. On the whole the rocks of this series are relatively impermeable and serve as an effective confining layer for the water that gets into the underlying Lower Cretaceous strata. In localities where they have been dropped down by faulting and lie against the Lower Cretaceous rocks they may serve as a barrier and cause the water to rise to the surface as springs. It appears that such structure is responsible for the large springs near Balmorhea.

The Upper Cretaceous series in this region has a thin bedded flaggy or hard platy limestone at the base, which weathers to a noticeable rusty-red and yellow color on exposure. When freshly broken, most of this limestone

is of chalky-white color, and most hillsides formed on it are covered by dense flaggy slabs that ring when struck with a hammer. Somewhat higher in the section these beds become more thinly stratified and shaly, and some of the massive limestone weathers into chips like a shale but retains in the flakes the firmness of a dense white limestone.

Above the limestones of Upper Cretaceous age near Balmorhea is a considerable thickness of clay, described as blue clay by drillers when it is fresh and below the zone of weathering. In some exposures the weathered surface is soft and of ochre-yellow color. This material being non-resistant to erosion, is eroded into featureless valleys where it is unprotected by a capping of harder rocks. It is usually covered in the valleys by an overwash of later detritus and on steep hillsides by talus and landslides from the hard rocks above.

Exposures of Upper Cretaceous strata that are relatively limited in area may be seen at several places between Balmorhea and the Davis Mountain front. The beds are most prominently exposed immediately beneath the lavas on the steep front of the Davis Mountains and in the foothills, and there are several extensive exposures in the broad anticlinal valley extending from Star Mountain to near Gomez Peak. One of the most complete exposures of the Upper Cretaceous series in normal sequence is found high on the slope on the east side of Gomez Peak under the steep escarpment, but even here the soft clay beds are much obscured by slides and talus of the hard rock from the ledges above. An exposure of these beds is found in the west half of section 15, block 13, H. and G. N. R. R. survey, a little more than a mile northeast of Phantom Lake, but this section is far from complete. A good exposure of the clay that overlies the limestone in the lower part of the Upper Cretaceous is found in a cut on the north bank of Toyah Creek about

1-1/2 miles west of Balmorhea.

The thickness of the Upper Cretaceous series has not been determined in the vicinity of Balmorhea. The section exposed on the steep front of Gomez Peak indicates that the series is probably at least 500 feet thick and it may be thicker beneath some of the valleys. Two wells approximately 400 feet deep, at Balmorhea, are reported to have penetrated clay and shale to the bottoms of the holes. A well 600 feet deep about 20 miles south-east of Balmorhea in the valley of Limpia Creek at the east end of the Barrilla Mountains is reported to have gone through blue clay, probably all of Upper Cretaceous age, all the way. However, the east end of these mountains has a synclinal structure and the Upper Cretaceous beds may be uncommonly thick in the area. The evidence suggests a thickness of at least 500 or 600 feet in most places, but the thickness may be considerably greater than 600 feet in other places.

Tertiary volcanic rocks.- After the deposition of the Cretaceous rocks in this region there was a withdrawal of the seas and slight folding of the previously deposited strata and the rocks were subjected to considerable erosion.

Volcanic rocks having a total thickness of 1,500 to 2,000 feet were laid down on the eroded surface of the Upper Cretaceous rocks during early Tertiary time. These rocks form the capping of the Davis and Barrilla Mountains, and of numerous ridges and hills on their flanks. They were probably originally spread far beyond their present area of outcrop, but have been removed from many parts of the area by erosion. The lowest bed of the volcanic succession is generally white tuff or ash, indicating that the first volcanic eruptions were of the explosive type. Above the basal tuffs is a widespread bed of volcanic breccia, composed of angular fragments of the volcanic rock in a matrix of lava. The breccia is overlain

by lava flows of rhyolitic composition. Above these are beds of tuff and at least two flows of lava which are of trachytic composition. The different parts of the series are so closely related that they are probably the product of a single cycle of volcanic activity.

Since the volcanic rocks form the summits of the Davis and Barrilla Mountains, they occupy a belt of rainfall greater than that of the adjoining lowlands. Much of the lava is exceedingly porous because it is fractured and jointed and some of it is slaggy and naturally full of cavities. It thus absorbs much of the water that falls on it. Some of the layers of tuff are clay-like and impermeable so that water descending through the overlying permeable lavas does not pass through them but flows along their surfaces to emerge in springs along their outcrops. In general the volcanic rock rests on the impervious clays of the Upper Cretaceous, and where this contact lies above the gradient of the main drainage channels, as it does in many places, most of the water absorbed by the volcanic rocks is again fed into the surface drainage or stream and terrace gravels within the mountain area. The volcanic series dips below stream gradients in certain structurally depressed areas, and it is likely that where this occurs the porous lava forms important reservoirs for the accumulation and storage of ground water below the level of the streams. The water thus stored is probably prevented from penetrating directly to the Lower Cretaceous limestone by the relatively impermeable basal volcanic tuff and Upper Cretaceous clay. In some localities, however, it may eventually reach the limestone by first rising into the streams or into the gravels underlying them and then moving down stream to structurally high areas in which the gravels lie directly on the limestone. The most important of these structural depressions near Balmorhea is on the axis of a long syncline extending from Limpia Creek where it flows between the Davis and Barrilla Mountains northwestward beyond the valley of Cherry Creek.

After the extrusion of the lava in the Davis Mountain region, and presumably while its surface was being eroded, the area was subjected to gentle deformation, which produced a series of broad folds involving the lavas and underlying rocks. The area of the volcanics seems to have been depressed somewhat, perhaps in partial compensation for the weight of the lava masses. This deformation took place in mid-Tertiary, and perhaps in Miocene time.

Late Tertiary, Pleistocene and Recent alluvial deposits.--Gravels and alluvial deposits overlie the bedrock in the lowlands northeast of the Davis Mountains. Available records of water well drilling in the Balmorhea area show that the gravels in a few places are 50 feet deep or slightly deeper but usually are from 15 to 25 feet deep. In the places the Cretaceous bedrock is exposed at the surface and the gravel is absent.

The relation of the Balmorhea springs to the geology

In the foregoing summary of the geology references have been made to the distribution of the rock formations about Balmorhea and to the geologic structures in which these formations are involved. The accompanying areal geologic map and profile showing the structure are drawn to the same vertical and horizontal scale, and if the interpretations are correct should give a true picture of the position and attitude of these formations on the surface and underground.

The profile section follows a line drawn from the triangulation monument on the prominent lava hills about 2 miles southwest of Phantom Lake Spring, in a northeasterly course approximately transverse to the main structural lines of the district.

The distribution of the rocks which nearly everywhere are covered by gravels and other surface formations is interpreted from the geologic mapping of the outcrops that are available and the structure indicated by that evidence. The thickness of the Upper and Lower Cretaceous formations is taken as 500 feet for each unit, which is probably within about 100 feet of being the correct thickness.

There is an unconformity between the Upper Cretaceous series and the overlying lavas, the Cretaceous series having been folded into gentle

anticlines and synclines and extensively eroded in the epoch between the withdrawal of the Upper Cretaceous sea and the deposition of the Tertiary lavas. Therefore the part of the Cretaceous section that will be found immediately beneath the base of the lavas in any particular locality cannot be predicted.

Both lava flows and Cretaceous strata were involved in folding subsequent to the distribution of the lava, but this is relatively insignificant in the small section of these beds included in the diagram. The faulting to which the whole section has been subjected, and which is an important feature in the section, is believed to be of early Pleistocene age and affected all of the rocks in the area except such Pleistocene gravels or alluvium as have accumulated since that deformation. Aside from the permeability of the strata themselves, the geologic structure is undoubtedly the controlling factor that determines the movement of the water underground and the location and existence of the big springs.

The principal area of intake of the water discharged by the springs apparently is in the anticlinal valley adjacent to the steep front of the Davis Mountains. (see p. 11 and fig. 1, p. 82) where a large part of the flow of the mountain streams disappears into Lower Cretaceous limestone or into stream gravels that overlie the limestone. The limestone dips to the northeast from the axis of the anticlinal valley to the axis of a major northwest-southeast trending syncline and then rises to the surface in the vicinity of Phantom Lake Spring. The axis of the syncline is marked by a range of volcanic hills, a part of which is shown in the southwestern corner of the geologic map (Plate 1 p. 81). In this syncline the base of the Lower Cretaceous section probably sinks to a depth of 1,000 to 1,200 feet or more below the major stream grades. The water probably fills the limestone in this structure to the height of its confining rim on the northeast, and is brought to or near the surface at the

outcrop of the reservoir beds in the vicinity of Phantom Lake. About 1,000 feet northeast of Phantom Lake there is a clearly defined fault on the south side of which the Lower Cretaceous strata have been tilted up steeply and on the north side of which the whole section has been dropped down, bringing relatively impermeable Upper Cretaceous strata against the Lower Cretaceous limestone. The Upper Cretaceous rocks on the north side of this fault may thus constitute an underground barrier to the passage of ground water at this place and cause the water to rise to the surface in Phantom Lake Spring. The spring emerges in a channel developed by solution along a fissure in the massive limestone that extends underground below the present water surface. Phantom Lake stands over a sink hole where the limestone has fallen in, but there is no evidence of faulting at the lake. The course of surface drainage below Phantom Lake is in approximately the same direction as the line of the structure section.

Immediately to the northeast of the Phantom Lake fault the strata rise gently toward the surface in the same manner as near Phantom Lake. This rise might have a general tendency to bring the water up again, or such part of it as sinks in this fault block, but it is again interrupted by a fault, the surface trace of which is obscured but is believed to lie almost a mile northeast of Phantom Lake. (See Pl. 1, p. 81).

The structure is shown in conspicuous outcrops in the channel of the stream from Phantom Lake about one and one-half miles northeast of the spring and about 1,000 feet north of the Old Spanish Trail highway. At this place the upper part of the Upper Cretaceous lies at the surface and is capped by a remnant of Tertiary lava. Therefore the area is structurally low.

No evidence of irregularities is known between this outcrop and the vicinity of Toyahvale and the San Solomon and Giffin springs. The broad

relatively level, alluvium-covered area that extends many miles both northwest and southeast from Toyah Creek in this part of the district is evidently eroded on the little-resistant clays of the upper part of the Upper Cretaceous section.

The valley within a radius of a mile of the San Solomon and Giffin springs is covered everywhere with a mantle of gravel or other detritus but a clue as to the probable character of the structure in their vicinity is afforded by the evidence on the south slope of the lava hills that project into the valley about a mile east of the springs. It is believed that the straight line which bounds the southwestern side of the mass of volcanic rocks that extends northwestward from the Barrilla Hills marks the trace of a fault. On the southwest side of this fault the Lower Cretaceous limestone is believed to be near the surface, and on the northeast side the Upper Cretaceous clays are again faulted down against the water-bearing limestone. The springs themselves are fair evidence of this. The inferred relations are shown by the structure section.

The basin between the fault near San Solomon Spring and the Brogada Hills, which is the part of the valley in which Balmorhea is situated, is mostly if not entirely underlain at shallow depth by the clay of the Upper Cretaceous section. There is evidence of this fact in a number of outcrops and this general structure is also indicated in the regional relations. In this part of the valley the top of the Lower Cretaceous rocks probably is about 500 feet below the surface. At the western foot of the Brogada Hills there is considerable disturbance of the beds and the fact that they are composed of Tertiary lava indicates that these hills are a down-faulted block along the southwest side of which impermeable Upper Cretaceous rocks are faulted against the Lower Cretaceous limestone. If this assumption is correct

the fault has produced a more or less effective barrier to the northeastward movement of underground water in the limestone in this area.

The extensive plains that lie east of the Brogada Hills are, so far as present evidence goes, developed on a broad structurally depressed area that begins at the west with the syncline in the lava shown in the Brogada Hills. There is a reasonable probability that the fault line bounding these hills on the southwest separates the Balmorhea basin from the ground-water basin east of the Brogada Hills. Evidence of this is seen in the fact that so much of the ground water of the upper basin has been brought to the surface above this line of displacement. The ground water in the lower or Saragosa basin is probably derived in part from the overflow from the Toyah Creek drainage system, in part from the northern flanks of the Barrilla Mountains, and possibly in large part from the Lower Cretaceous limestone near the center of the Barrilla Mountains where it may be close enough to the surface to receive water through the gravels from surface streams.

The blanket of gravels that underlies the Saragosa plains is thicker than the gravel deposits in the Balmorhea basin. According to general reports of water wells that have been drilled, it seems that this thickening takes place abruptly near the settlement of Brogada and increases eastward. The Saragosa Basin is filled with a coarse impermeable boulder conglomerate. Water wells sunk through this upper gravel find water in a loose and clean sand below it. Tests of this sand indicate that, in general, it does not yield enough water to warrant considering it as a possible source of important new irrigation supplies.

According to the foregoing interpretations of the structure the Lower Cretaceous limestones in the Saragosa area should yield artesian water if they are as porous or channelled as they are in many places. These beds are structurally lower at Saragosa than they are either to the east or west of that basin, as they lie in the southern extension of the Toyah trough. There is

little to indicate the depth at which the limestones would be encountered in this area, but it seems unlikely that they would be deeper than 1,000 or 1,200 feet. It is probable that the ground water in this trough is derived largely from areas other than those that supply the springs near Balmorhea, and the Lower Cretaceous limestone in the Saragosa basin east of Brogada may be, therefore, a promising source of additional water supply.

Occurrence of ground water in the Balmorhea area

Springs

The locations of the Balmorhea springs are shown on plate 1, p. 81, and figures 1 and 2. pp 82-83. The springs are of two classes, artesian springs and gravity springs. The San Solomon, Phantom Lake and Giffin springs belong to the first class, and the Saragosa, West Sandia, East Sandia and Toyah Creek springs to the second. In the preceding sections of this report, Phantom Lake, San Solomon and Giffin springs have been called the "large springs". The artesian springs are those whose waters issue under artesian pressure and the gravity springs are those whose waters flow because the water table is above the land surface at the site of the spring and the water therefore tends to drain from the water-bearing beds.

Artesian springs.- The artesian character of Phantom Lake, San Solomon and Giffin springs is proved by the fact that they appear at the surface of the ground, or would flow at the surface, if they were permitted to do so. Phantom Lake Spring issues from an A-shaped opening in the side of a hill of Lower Cretaceous limestone. It is located at the northwestern edge of a large depression, formed, it appears, as the result of the collapse of the roof of a cavern in the limestone. The water flows into a small pond, called Phantom Lake, and thence is carried through a canal to irrigated lands in the valley below.

San Solomon and Giffin springs rise from gravels and originally came to the surface, forming a swamp of considerable size, which drained into Toyah Creek. The San Solomon Spring now issues from gravels in the bottom of a large well-built bathing pool. The water is beautifully clear and the discharge at all times has the magnitude of a good sized creek. It is not surprising, therefore, that the spring is one of the leading scenic attractions of the region. The Giffin Springs rise from gravels at or near the head of

four short pits or trenches that finger out from a central outlet trench. When first used for irrigation the flow of the springs was diverted into canals that were cut only a few feet below the surface. Eighteen or twenty years ago, with the hope of obtaining a large increase in flow by lowering the points of discharge of the springs, trenches and pits of considerable depth were excavated at both sites. Then new and deeper outlet canals were constructed, the one at San Solomon Spring having a depth of about 20 feet and that at Giffin Spring a somewhat lesser depth.

The high and low-level canals at both springs are available for diversion, but the low-level canal from San Solomon Spring is used infrequently because its use does not increase the flow sufficiently to justify the inconvenience of distributing water from the lower level.

In the course of the investigation, efforts were made to determine the flow of San Solomon Spring when discharging into the high-level canal and when discharging into the low-level canal, but the demands for water for irrigation would not permit an accurate determination of the flow at the respective levels. However, a few measurements were made. On November 10, 1931 the discharge through the high-level canal was found to be 30.9 second-feet and in three measurements on November 20 and 21, ten and eleven days later, the discharge through the low-level canal was 37.7, 37.6 and 38.0 second-feet. These measurements indicate a difference in flow between the two levels of six to seven second-feet. The spring had been discharging through the high-level canal for about a month prior to the measurement on November 10, and during that time the flow had not varied more than one to two second-feet. It had been discharging through the low-level canal for eight or nine days before the measurements at that canal were made. The computed difference in flow at the two levels, therefore, may not have been far from the correct difference

for the period. Whether or not the flow would gradually decrease if the outlet should be maintained at the lower level for a long time is not known. It appears probable that it might.

The discharge of all three springs was well sustained even during several successive dry years. Phantom Lake Springs has a somewhat wider variation in flow than the other two. The lowest discharge of San Solomon Spring recorded by the Geological Survey, was 26.5 second-feet on April 26, 1923 (p. 55) and the highest was about 71 second-feet on October 7, 1932 (p. 66). The lowest recorded flow of Phantom Lake Spring was 10 second-feet, October 16, 1931, and the highest 114 second-feet October 2 and 3, 1932 (p. 62). The discharge of the Giffin Springs is relatively small, the smallest recorded flow being 2.9 second-feet on March 4, 1925 and the largest between 6 and 7 second-feet in October 1932. Daily records of the discharge of the springs during a period of about two years in 1931 to 1933 are given on the following pages: Phantom Lake Spring (pp. 61-63), San Solomon Spring (pp. 65-67), Giffin Spring (pp. 63-64).

The flow of Phantom Lake Spring and San Solomon Spring increases very soon after heavy rains, the rise occurring more quickly and the rate of increase being greater in Phantom Lake Spring than in San Solomon. For example: The discharge of Phantom Lake Spring had remained relatively constant at about 13 second-feet from December 21, 1931 to February 22, 1932. On February 20 to 22, 1932, 2.41 inches of rain was recorded at Balmorhea, the heaviest fall being on February 21, and as a result all the canyons had a substantial flow of storm water. The discharge of Phantom Lake Spring began to increase on February 22 and reached 21 second-feet on February 23,

The flow of San Solomon Spring which had remained constant about 30 second-feet for several weeks began to increase on the afternoon of February 23, about 24 hours after the rise started at Phantom Lake. The flow continued to increase gradually and reached a maximum of 36 second-feet on February 27 after which it gradually decreased.

The rainfall in this district from August 30 to September 30, 1932 amounted to 14.36 inches and was the heaviest on record. The first heavy rain, 2.72 inches, occurred August 30, and was quickly followed by large runs of storm water from the mountain canyons. The flow of Phantom Lake Spring which had been relatively constant at 12 to 13 second-feet for several months, began to increase on the evening of August 29, and reached about 46 second-feet on September 11. The flow then gradually decreased to about 25 second-feet on September 26, when it again began to increase and reached a maximum of 104 second-feet on October 3. The flow of San Solomon Spring, which had remained at 32 to 33 second-feet for several months, began to increase on August 29 and continued to increase until September 17, when it was about 63 second-feet. The discharge then slowly decreased reaching a minimum of about 54 second-feet on September 23 when it again started to increase and reached a maximum of about 73 second-feet on October 7.

The temperature of Phantom Lake, San Solomon and Giffin springs normally is high. The temperature of all three was the same (78° F.) in October 1931. After the increase in the flow of the springs in September 1932 the temperature was 71° in Phantom Lake Spring, and 74° in San Solomon Spring. On March 11, 1933 the temperature was 77-1/2° in Phantom Lake Spring and 73-1/2° in San Solomon Spring.

Ordinarily the discharge from the springs is perfectly clear, but very soon after the start of the abrupt increases in flow in August and September 1932, the water from Phantom Lake and San Solomon springs became cloudy due

to the suspension of a very fine light colored sediment in the water.

Normally the mineral content of the water of the springs is rather high but the water becomes quite fresh during periods of peak discharge. This is well illustrated by the table on page 27 which gives the comparative results of analyses of samples obtained during periods of moderate flow in the fall of 1930 and of high flow in the fall of 1932. When the samples were taken in 1930 the discharge was not measured but was known to be at moderate or rather low stage. In the fall of 1932 the discharge of all the springs was abnormally high. When they were sampled on September 12 and 13, 1932, Phantom Lake Spring had a discharge of 45 second-feet and San Solomon Spring 64 second-feet. When sampled on October 7, 1932 Phantom Lake Spring had a discharge of about 82 second-feet, San Solomon Spring 71 second-feet and Giffin Spring 6 second-feet. The results show that the decline in the mineral content of the water at high stages is remarkably large. The decline in the total dissolved solids was as follows:

Total dissolved solids in waters from the large springs
near Balmorhea at different dates

Phantom Lake Spring		San Solomon Spring		Giffin Spring	
Date	Total dissolved solids (parts per million)	Date	Total dissolved solids (parts per million)	Date	Total dissolved solids (parts per million)
Oct. 28, 1930	2,309	Oct. 28, 1930	2,196	Dec. 6, 1930	2,098
Sept. 12, 1932	723	Sept. 13, 1932	875		
Oct. 7, 1932	144	Oct. 7, 1932	562	Oct. 7, 1932	547

The most outstanding feature of this record is the change in the discharge at Phantom Lake Spring from a highly mineralized water to water containing probably very little more mineral matter than the Davis Mountain streams. The decline in the mineralization of the water from San Solomon and Giffin Springs, while large, was markedly less than the decline in Phantom Lake Spring.

Analyses of water from the large springs near

Balmorhea, Texas

(Parts per million)

Analyzed by M. D. Foster and L. A. Shinn

Spring	Date of Collection	Total dissolved solids	Cal- cium (Ca)	Magne- sium (Mg)	Sodium and Potassium (Na / K) (calc.)	Bicar- bonate (HCO ₃)	Sul- phate (SO ₄)	Chlo- ride (Cl)	Ni- trate (NO ₃)	Iron (Fe)	Silica (SiO ₂)	Total hardness as CaCO ₃
Phantom Lake	Oct. 28, 1930	2,309	191	86	473	285	691	655	0.55	0.04	19	830
do.	Sept. 12, 1932	<u>a/</u> 723	81	27	<u>a/</u> 139	170	905	186	1.4	-	-	<u>a/</u> 313
do.	Oct. 7, 1932	<u>a/</u> 144	<u>b/</u> 44	-	<u>a/</u> 10	131	<u>b/</u> 20	4	-	-	-	111
San Solomon	Oct. 28, 1930	2,196	190	80	448	286	651	610	0.90	0.04	19	803
do.	Sept. 13, 1932	<u>a/</u> 975	102	35	<u>a/</u> 200	264	270	238	0.38	-	-	<u>a/</u> 398
do.	Oct. 7, 1932	<u>a/</u> 562	<u>b/</u> 88	-	<u>a/</u> 93	189	<u>b/</u> 175	100	-	-	-	276
Giffin	Dec. 6, 1930	<u>a/</u> 2,098	189	80	<u>a/</u> 437	284	635	608	-	-	-	<u>a/</u> 800
do.	Oct. 7, 1932	<u>a/</u> 547	<u>b/</u> 80	-	<u>a/</u> 109	183	<u>b/</u> 140	124	-	-	-	234

a/ Calculated.

b/ By turbidity.

The persistent discharge of large quantities of water by the large springs during months or even years of drought indicates that they are fed from a large reservoir or from a series of reservoirs. The water must be derived in large part from rainfall and run-off that enters the reservoir at no great distance from the springs. This is indicated by the abrupt increase in the discharge of Phantom Lake and San Solomon springs following heavy rains and large runs of muddy water from the mountain canyons, by the decrease in the dissolved minerals in the spring water, and by the lowering of the temperature of the water and the cloudy character of the water at such times. The data all tend to substantiate the conclusions in the section on the geology and the relation of the springs to the geologic structure. Analyses of the water from the large springs are given on page 27.

Gravity Springs.- Saragosa, West Sandia, and East Sandia springs rise from gravels in the bottom of shallow drainage channels near Balmorhea. The Toyah Creek springs issue at numerous points in the bed of Toyah Creek in a stretch of three or four miles southwest and northeast of Balmorhea. The uppermost point of emergence of the Saragosa and Toyah Creek springs shifts materially up and down stream from year to year, and season to season. The position of the principal West Sandia Spring now appears to be fixed at a point in a shallow drainage channel on the southeastern border of Balmorhea, but it is reported that about 30 years ago this spring emerged more than half a mile southwest of Balmorhea, or about a mile up the valley from its present location. For a time after the heavy rains of August and September 1932, small seeps in the channel above and below the spring were discharging more ground water than the spring itself. The East Sandia Spring appears at the base of the western slope of the Brogada hills, and so far as could be learned it has always been where it is today.

As determined by instrumental levelling the heads of the Saragosa and Toyah Creek springs in November 1931 were a few feet below the water levels in nearby shallow wells that draw from the gravel; likewise, the outlet of the main West Sandia Spring was one to two feet lower than the water level in well 20, one of the shallow observation wells that are discussed later in this report. The outlets of the East Sandia Spring apparently are about at the level of the shallow water table immediately above the spring, but this was not exactly determined.

Saragosa Spring enters Toyah Creek and joins the ground water discharge of that stream a few hundred feet above the Balmorhea bridge. The measured discharge of the spring in second-feet was as follows: November 6, 1932, 9.2; January 23, 1933, 6.7; March 14, 1933, 6.7; May 16, 1933, 6.4; July 11, 1933, 5.6.

The inflow of ground water into Toyah Creek down to the Moore dam about a mile below the Balmorhea bridge, consisting of the combined flow of the Saragosa and Toyah Creek springs, was as follows: November 6, 1932, 30.4 second-feet; January 23, 1933, 12.8 second-feet; March 14, 1933, 9.4 second-feet; May 16, 1933, 8.9 second-feet; July 11, 1933, 8.6 second-feet. According to records of the Reeves County Water Improvement District the combined discharge of the springs down to the same point ranged from 4.8 to 6 second-feet during December 1931 and January 1932 and from 5.8 to 8.3 second-feet in June, July and August 1932. The rates of flow, shown by the measurements obtained prior to the floods in August and September 1932, probably are much nearer the average long time flow than those obtained during the months succeeding the floods.

The daily discharge of West Sandia Spring from November 17, 1931 to August 28, 1933 is given in the tables on pp. 72 and 73. According to one set of measurements on October 17, 1932 shortly after the heavy rains the

discharge of the principal spring amounted to .6 second-foot, and the seepage inflow above it to .2 second-foot while the inflow between the spring and the gaging station 4,000 feet below it amounted to 1.4 second-feet. A part of this was diverted above the gage and was not taken into account in the gaging records. The mean monthly discharge of East Sandia Spring in second-feet and the monthly run-off in acre-feet from November 23, 1931 to September 23, 1933 is given in the table on page 74.

The temperature of the water from Saragosa and East and West Sandia springs is materially lower than the temperatures of the waters of the large springs. On March 14, 1933 the following temperatures were recorded: Saragosa Spring, 67-1/2 degrees F.; West Sandia Spring 63-1/2°; East Sandia Spring, 3 outlets, 55 degrees, 58 degrees and 64 degrees.

The Saragosa, West Sandia, East Sandia, and Toyah Creek springs quite clearly are gravity springs and have their source in the gravels. They appear at levels slightly below the water table in the gravels. Their temperatures are much lower than those of the large springs and are not materially different from the temperatures of the water from shallow wells in the vicinity. The discharge of the Saragosa and Toyah Creek springs apparently increases and decreases and the position of the uppermost outlets shifts up and down the streams with the rise and fall of the water table. The position of East Sandia Spring is stationary and that of the main West Sandia Spring is nearly stationary. This may be due to the fact that they are fed from solution channels in the cemented gravels, and that these channels are not directly connected with the shallow water table. The basal conglomerate is encountered in nearly all the shallow wells in the Balmorhea district and is known to be cavernous in places. Irrigation streams amounting to a second-foot or more have been known to disappear into these passages.

Shallow ground water.- Most of the water used for domestic and stock purposes in this area is obtained from wells. A description of forty-one of the wells is given in the table on pages 32-33 and the location of the wells is shown in figure 2, ^{p.83/} each well being given a number that corresponds to the number assigned to it in the table. In the vicinity of Balmorhea, all the wells terminate in the gravels, none are more than 60 feet deep and most of them are less than 30 feet deep. The depth to water was measured in all the wells and the measuring points at the top of most of them were connected by lines of levels and the elevations of these points referred to a common assumed datum. In this way the water table map, figure 2, ^{p.83/} was prepared. This map indicates that the slope of the water table is in the same direction as that of the land and that the movement of the ground water in the gravels, with some variations, is directly down the valley. It shows further that the elevations of the Saragosa and West and East Sandia springs conform closely with the elevations of the water table in their vicinities.

Thirteen wells were selected for observation and measured periodically over a period of several years. The records of these measurements are shown in the tables on pp. 34-40. These records show that the water levels in the wells are in general moderately low, but that they rise when heavy rains occur, when Toyah Creek is in flood or when nearby lands are irrigated. It is evident, therefore, that the principal source of the ground water in the gravels is rainfall and seepage from irrigation canals and irrigated lands. Seepage from the large springs doubtless contributes to the shallow ground water. As previously stated the water in the gravels is believed to be the source of the Saragosa, West Sandia, East Sandia and Toyah Creek springs.

Records of wells in vicinity of Balmorhea, Texas

No.	Distance from Balmorhea (miles)	Owner	Depth (feet)	Water level		Alti- tude of B.M. <u>a/</u>	Use of water <u>b/</u>	Method of lift <u>c/</u>
				Below B. M. (feet)	Date			
1	3 $\frac{3}{4}$	southwest	Balmorhea Livestock Company	175	16.29 Nov. 10, 1931	3349	D,S	W
2	2	southwest	A. C. Schreger	14	13.99 Oct. 31, 1930	3286	D,S	W
3	1 $\frac{3}{4}$	southwest	Charles Weinacht	15	13.73 do.	3275	S	W
4		do.	A. W. Wigley	20	14.39 do.	3278	D,S	W
5	1 $\frac{3}{4}$	south-southwest	Humble Oil & Refining Company	14	14.47 Nov. 13, 1931	3272	D,S	H
6	1 $\frac{3}{4}$	south-southwest	S. Garcia	20	13.91 Oct. 31, 1930	3266	D,S	H
7	2 $\frac{3}{4}$	south-southwest	S. A. Sharpe	20 $\frac{1}{2}$	12.86 do.	3286	D,S	H
8	1 $\frac{3}{4}$	south-southwest	H. A. Jones	20 $\frac{1}{2}$	14.98 do.	3263	D,S	W
9	1 $\frac{3}{4}$	southwest	do.	19	11.65 Nov. 13, 1931	3265	S	H
10		do.	W. A. Knapp	17	12.22 Nov. 1, 1930	3264	N	E
11	1	southwest	Mrs. Nell B. Westerman	20	11.61 do.	3251	S	W
12	3 $\frac{3}{4}$	northwest	-- Hill	22	11.70 Nov. 3, 1930	3234	D,S	H
13	1 $\frac{3}{4}$	west	Rosenbaum Company	15	13.45 Nov. 1, 1930	3231	D,S	W
14	1 $\frac{3}{4}$	southwest	T. A. Odell	10	7.95 do.	3228	-	H
15	1 $\frac{3}{4}$	south	Rosenbaum Company	9	9.35 do.	3234	D,S	H
16		do.	A. W. Wigley	17	6.89 do.	3225	N	None
17		do.	Rosenbaum Company	11	9.86 do.	3233	D,S	H
18	1	southeast	Mrs. Paul Renz	12	4.20 Nov. 13, 1931	3221	S	W
18A	1 $\frac{1}{2}$	southeast	Unknown	20 $\frac{1}{2}$	4.71 do.	3211	Test	hole
19	In Balmorhea	do.	Unknown	5.33	Nov. 6, 1931	3221	N	None
20	do.	Toyah Valley State Bank	7	7.99	Nov. 1, 1930	3214	N	None
21	1 $\frac{1}{2}$	northeast	J. F. Meier	24	23.27 do.	3208	D,S	W
22	1	northeast	P. V. & S Ry.	10	5.56 do.	3198	N	None
23		do.	Unknown	Unknown	2.16 Oct. 31, 1931	3194	N	None
24	1 $\frac{1}{4}$	northeast	E. P. Stuckler	13	11.68 Nov. 6, 1930	3186	D,S	W
25		do.	Mike Tesero	21.5	10.63 Nov. 1, 1930	3188	D,S	H
26	1	north-northeast	J. P. Cole	17	14.65 Nov. 3, 1930	3198	D,S	H
27	1 $\frac{1}{4}$	northeast	R. W. Vanderson	20	20.66 do.	3190	D,S	H
28	1 $\frac{1}{4}$	north-northeast	E. W. Backus	18	16.60 Nov. 1, 1930	3197	D,S	W
29	1 $\frac{1}{2}$	north	Perry Wegnon	50	34.35 Nov. 3, 1930	3204	D,S	W
30	1	northwest	W. E. Gould	36	26.88 Nov. 1, 1930	--	D,S	W

Records of wells in vicinity of Balmorhea, Texas -- Continued

No.	Distance from Balmorhea (miles)	Owner	Depth (feet)	Water level		Altitude of B.M. a/	Use of water b/	Method of lift c/
				Below B. M. (feet)	Date			
31	1½	northwest	W. E. Gould	58	30 Nov. 3, 1930	-	P	E
32	do.	do.	do.	53	41.51 Nov. 1, 1930	-	N	None
33	do.	do.	do.	60	46.82 do.	-	P	G
34	1¾	north-northeast	Mrs. W. E. Gould	19	16.52 Nov. 3, 1930	3188	N	W
35	3¾	east-northeast	State of Texas	220	180.4 Nov. 5, 1930	-	D, S	W
36	4	east-northeast	A. H. Mills	117	112.9 do.	-	N	W
37	6	northeast	Saragoso School	158	139.2 Nov. 3, 1930	-	D	W
38	6¾	northeast	J. H. Downs	158	137.6 do.	-	D, S	W
39	¾	east	G. F. Renz	26	9.80 Nov. 2, 1931	3210	D, S	C
40	do.	do.	do.	18	9.21 do.	3209	D, S	S
41	In Balmorhea	Reeves County Water Improvement Dist.		202	5.56 Nov. 1, 1930	-	-	None

a/ Assumed datum.

b/ D, domestic; S, stock; P, public supply; N, not used.

c/ W, windmill; H, hand; E, electric motor; G, gas engine; S, siphon.

Well measurement records

(For description of wells see preceding table)

3. Charles Weinacht, 1 $\frac{1}{2}$ miles southwest of Balmorhea postoffice.
 Measuring point, top of edge of pipe clamp, south side, at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Oct. 31, 1930	13.73	Dec. 10, 1932	7.77	May 19, 1933	<u>e/</u> 8.74
Mar. 8, 1931	14.50	Dec. 14	7.95	May 20	9.00
Dec. 11	13.72	Jan. 21, 1933	7.59	May 27	8.72
Jan. 19, 1932	<u>a/</u> 14.26	Jan. 28	8.12	June 2	8.40
Feb. 15	15.00	Feb. 4	8.16	June 10	8.92
Apr. 13	<u>b/</u> 9.74	Feb. 11	8.18	June 17	<u>e/</u> 7.60
May 30	11.61	Feb. 18	8.30	June 24	<u>e/</u> 6.95
Aug. 4	11.52	Feb. 25	8.43	July 8	<u>f/</u> 8.47
Aug. 21	11.30	Mar. 4	<u>d/</u> 8.75	July 15	9.65
Sept. 17	7.60	Mar. 4	9.15	July 27	<u>f/</u> 8.85
Oct. 1	<u>c/</u> 3.56	Mar. 11	<u>e/</u> 4.71	Aug. 5	8.68
Oct. 15	5.96	Apr. 1	8.35	Aug. 19	9.54
Oct. 22	6.18	Apr. 9	8.50	Aug. 26	11.17
Oct. 30	6.58	Apr. 16	<u>e/</u> 8.20	Sept. 2	<u>g/</u> 11.08
Nov. 5	6.61	Apr. 23	8.40	Sept. 23	9.49
Nov. 12	6.91	Apr. 29	8.45		
Nov. 19	5.49	May 7	9.00		

5. Humble Oil and Refining Company, 1 $\frac{3}{4}$ miles south-southwest of
 Balmorhea postoffice.
 Measuring point, Benchmark in top of wood cover, 2 feet above ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 13, 1931	14.47	Dec. 10, 1932	7.19	May 14, 1933	9.90
Dec. 11	11.42	Jan. 10, 1933	7.68	May 20	10.15
Jan. 19, 1932	<u>a/</u> 12.86	Jan. 14	7.86	May 27	10.42
Feb. 15	<u>b/</u> 11.04	Jan. 28	8.22	June 3	10.65
May 30	10.08	Feb. 4	8.40	June 10	11.03
Aug. 4	11.11	Feb. 11	8.60	June 17	11.22
Sept. 17	11.50	Feb. 18	8.65	June 24	11.80
Oct. 1	<u>c/</u> 8.61	Feb. 25	<u>d/</u> 8.70	July 8	12.68
Oct. 15	6.79	Mar. 11	9.37	July 15	13.01
Oct. 22	6.72	Apr. 1	9.67	July 29	13.32
Oct. 30	6.54	Apr. 9	9.30	Aug. 5	13.52
Nov. 5	6.77	Apr. 16	9.37	Aug. 26	13.95
Nov. 12	6.92	Apr. 23	9.58	Sept. 2	<u>g/</u> 14.10
Nov. 19	7.01	Apr. 26	9.62	Sept. 23	13.66
Dec. 3	7.24	May 7	9.76		

7. S. A. Sharpe, $2\frac{3}{4}$ miles south-southwest of Balmorhea postoffice.
Measuring point, benchmark in plank, north side, at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Oct. 31, 1930	12.86	Jan. 19, 1932	9.44	Aug. 4, 1932	15.73
Mar. 8, 1931	<u>b/</u> 10.37	Feb. 15	<u>a/</u> 8.00	Aug. 20	16.43
Nov. 7	15.08	Apr. 13	9.50		
Dec. 11	10.96	May 30	12.89		

10. W. A. Knapp, $1\frac{1}{2}$ miles southwest of Balmorhea postoffice.
Measuring point, benchmark in east end and south side of middle cross
beam, 1 foot above ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 1, 1930	12.22	Dec. 12, 1932	6.08	May 14, 1933	8.80
May 8, 1931	11.00	Jan. 10, 1933	7.13	May 20	9.05
Nov. 4	9.89	Jan. 14	7.47	May 27	9.03
Dec. 11	13.02	Jan. 21	7.30	June 3	8.70
Jan. 19, 1932	<u>a/</u> 12.79	Jan. 28	7.45	June 10	9.18
Feb. 13	<u>b/</u> 11.00	Feb. 4	7.60	June 17	8.80
Apr. 13	11.00	Feb. 11	7.98	June 24	9.12
May 30	10.8	Feb. 18	8.16	July 8	9.22
Aug. 4	10.85	Feb. 25	8.35	July 15	9.78
Aug. 20	11.59	Mar. 4	<u>d/</u> 8.63	July 29	9.69
Sept. 17	6.60	Mar. 11	<u>f/</u> 8.22	Aug. 5	9.98
Oct. 1	<u>c/</u> 4.51	Apr. 1	8.30	Aug. 12	10.52
Oct. 3	4.30	Apr. 9	8.78	Aug. 19	10.18
Oct. 15	5.18	Apr. 16	8.23	Aug. 26	10.70
Oct. 22	5.38	Apr. 23	8.64	Sept. 1	10.78
Oct. 30	5.31	Apr. 29	8.40	Sept. 23	<u>g/</u> 7.58
Nov. 5	6.36	May 7	8.35		

12. -- Hill, $\frac{1}{2}$ mile northwest of Balmorhea postoffice.
Measuring point, top edge of upper wood frame, east side opposite
cut in frame, at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 3, 1930	11.70	Jan. 21, 1933	8.03	May 23, 1933	8.85
Dec. 11, 1931	11.89	Jan. 28	8.16	June 4	9.13
Jan. 19, 1932	<u>a/</u> 13.10	Feb. 4	8.60	June 11	8.90
Feb. 15	13.00	Feb. 11	8.62	June 13	9.20
Apr. 13	<u>b/</u> 9.23	Feb. 14	8.70	June 25	9.45
May 30	11.42	Feb. 25	8.76	July 8	9.05
Aug. 4	9.90	Mar. 4	<u>d/</u> 9.00	July 15	9.08
Aug. 20	10.72	Mar. 11	8.36	July 29	9.50
Sept. 17	5.79	Apr. 1	8.63	Aug. 5	9.82
Oct. 15	<u>c/</u> 5.46	Apr. 9	<u>f/</u> 8.22	Aug. 11	<u>f/</u> 9.64
Oct. 22	5.79	Apr. 16	<u>f/</u> 8.15	Aug. 19	10.08
Oct. 30	6.18	Apr. 23	8.63	Sept. 2	<u>g/</u> 9.32
Nov. 5	6.46	Apr. 29	8.47	Sept. 23	8.66
Dec. 10	7.06	May 7	8.22		
Jan. 10, 1933	7.86	May 14	8.75		
Jan. 14	7.93	May 20	8.25		

16. A. W. Wigley, $\frac{1}{2}$ -mile south of Balmorhea postoffice
Measuring point, southwest edge of concrete cross-beam, opposite cut
in concrete at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 1, 1930	6.89	Mar. 21, 1932	<u>e/</u> 2.55	July 23, 1932	4.07
Mar. 8, 1931	4.20	Mar. 26	4.11	July 30	5.29
Nov. 12	4.91	Apr. 2	4.45	Aug. 6	5.86
Nov. 19	5.50	Apr. 9	4.49	Aug. 13	6.04
Nov. 23	5.73	Apr. 12	3.25	Aug. 16	6.16
Dec. 3	5.99	Apr. 16	4.13	Aug. 20	<u>e/</u> 5.47
Dec. 10	5.69	Apr. 23	4.41	Aug. 30	<u>c/</u> 5.35
Dec. 19	5.87	Apr. 27	<u>f/</u> 5.25	Sept. 3	4.25
Dec. 26	5.79	Apr. 30	3.87	Sept. 4	4.25
Dec. 29	5.83	May 7	4.59	Sept. 4	1.47
Jan. 2, 1932	<u>a/</u> 5.77	May 14	4.85	Sept. 4	<u>c/</u> .82
Jan. 9	5.67	May 28	4.95	Sept. 10	1.97
Jan. 16	5.76	June 3	<u>f/</u> 5.45	Sept. 17	2.76
Jan. 23	5.73	June 4	3.86	Sept. 24	2.91
Jan. 28	5.67	June 8	4.50	Oct. 1	<u>c/</u> 1.03
Jan. 30	4.79	June 10	<u>f/</u> 3.15	Oct. 9	2.37
Feb. 6	4.93	June 11	3.68	Oct. 15	2.17
Feb. 11	<u>c/</u> 2.35	June 17	4.05	Oct. 22	1.39
Feb. 13	4.15	June 18	<u>f/</u> 3.72	Oct. 29	2.15
Feb. 20	4.19	June 25	4.24	Nov. 5	2.73
Feb. 23	<u>b/</u> 2.55	June 30	4.66	July 29, 1933	4.45
Feb. 27	5.41	July 2	5.45	Aug. 5	<u>f/</u> 4.31
Feb. 29	3.53	July 4	<u>f/</u> 4.05	Aug. 26	4.13
Mar. 5	3.71	July 9	5.35	Sept. 2	<u>f/</u> 3.23
Mar. 12	4.22	July 16	5.56	Sept. 23	3.65
Mar. 19	4.21	July 22	<u>f/</u> 3.35		

20. Toyah Valley State Bank, in Balmorhea
Measuring point, cross in top of west brick in top row,
at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 1, 1930	5.69	Dec. 10, 1932	4.28	May 14, 1933	4.03
Mar. 8, 1931	5.70	Jan. 10, 1933	3.93	May 20	4.20
Nov. 10	5.67	Jan. 21	4.09	May 27	4.58
Dec. 11	5.60	Jan. 28	4.00	June 4	4.35
Jan. 19, 1932	a/ 5.43	Feb. 4	4.08	June 11	4.50
Feb. 15	b/ 3.95	Feb. 11	4.03	June 18	4.40
Aug. 4	4.9	Feb. 18	4.10	June 25	4.30
Aug. 20	4.95	Feb. 25	4.00	July 8	4.76
Sept. 17	3.8	Mar. 4	d/ 3.92	July 13	5.00
Oct. 1	c/ 2.56	Mar. 11	4.02	July 21	4.80
Oct. 15	3.51	Apr. 1	3.73	Aug. 5	4.79
Oct. 22	3.55	Apr. 9	3.74	Aug. 11	4.92
Oct. 30	3.96	Apr. 16	3.80	Aug. 19	4.62
Nov. 5	4.04	Apr. 22	3.67	Aug. 25	g/ 4.72
Nov. 12	4.14	Apr. 26	4.03	Sept. 2	4.04
Nov. 19	4.19	May 7	3.96	Sept. 25	3.36

22. On P. V. S. Ry. right-of-way, 1 mile northeast of Balmorhea postoffice.
Measuring point, benchmark copper nail on south side well curb, 3 feet
above ground level.

Date	water level above (+) or below (-) measuring point (feet)	Date	water level above (+) or below (-) measuring point (feet)	Date	water level above (+) or below (-) measuring point (feet)
Nov. 1, 1930	-5.56	Oct. 15, 1932	/ 1.92	Apr. 29, 1933	-1.18
Mar. 8, 1931	-4.50	Oct. 22	/ 3.22	May 7	-1.30
Dec. 10	-1.65	Oct. 20	/ 0.15	May 14	-1.32
Feb. 15, 1932	b/ -1.00	Nov. 5	- 0.04	May 20	-1.52
Apr. 13	-1.00	Nov. 10	- 1.28	May 23	-1.65
May 30	-0.4	Nov. 19	- 0.49	June 11	-1.78
Aug. 4	-1.13	Apr. 1, 1933	-1.00	June 11	-1.95
Aug. 20	-1.80	Apr. 9	-1.10	June 17	-2.00
Sept. 17	c/ /1.3	Apr. 16	-1.14	June 25	-2.00
Oct. 1	h/ /3.9	Apr. 23	-1.20		

23. Along east side of P. V. S. Ry. right-of-way 1 mile northeast of Balmorhea postoffice. Measuring point, benchmark in top of board lining, west side of well, about 2 feet below ground level.

Date	water level above (✓) or below (-) measuring point. (feet)	Date	water level above (✓) or below (-) measuring point. (feet)	Date	water level above (✓) or below (-) measuring point. (feet)
Oct. 31, 1931	-2.16	Nov. 5, 1932	-0.04	Mar. 11, 1933	-0.83
Dec. 10	-1.65	Dec. 10	-0.69	July 8	-2.19
Feb. 15, 1932	<u>b/</u> -1.00	Jan. 10, 1933	-0.28	July 15	-2.34
May 30	-0.04	Jan. 14	-0.38	July 22	-2.38
Aug. 4	-1.18	Jan. 21	-0.55	July 29	-2.50
Aug. 25	-1.8	Jan. 28	-0.58	Aug. 5	-2.52
Sept. 17	<u>c/</u> 1.3	Feb. 4	-0.52	Aug. 19	-2.72
Oct. 1	<u>h/</u>	Feb. 11	-0.54	Aug. 28	<u>g/</u> -2.82
Oct. 15	1.92	Feb. 18	-0.62	Sept. 2	-0.20
Oct. 22	3.22	Feb. 25	-0.65		
Oct. 30	0.05	Mar. 4	-0.72		

26. J. P. Cole, 1 mile north-northeast of Balmorhea postoffice. Measuring point, edge of concrete collar, northeast side of well, opposite cut in collar, 1 foot above ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 3, 1930	14.65	Jan. 14, 1933	11.65	May 20, 1933	12.42
Mar. 8, 1931	<u>c/</u> 12.16	Jan. 21	11.84	June 4	12.65
Nov. 3	14.44	Jan. 28	12.01	June 11	12.68
Dec. 11	14.95	Feb. 4	12.00	June 18	<u>e/</u> 12.23
Jan. 19, 1932	<u>a/</u> 15.96	Feb. 11	12.30	June 25	12.55
Feb. 15	<u>b/</u> 14.95	Feb. 18	11.28	July 8	12.59
Apr. 3	14.95	Feb. 25	<u>d/</u> 11.60	July 15	<u>e/</u> 12.21
May 30	14.4	Mar. 4	15.13	July 29	12.83
Aug. 4	14.45	Mar. 11	<u>e/</u> 11.57	Aug. 5	12.86
Aug. 20	14.47	Apr. 1	12.02	Aug. 11	12.55
Sept. 17	12.52	Apr. 9	12.14	Aug. 19	12.98
Oct. 15	<u>c/</u> 10.21	Apr. 16	12.23	Aug. 26	<u>g/</u> 13.22
Oct. 22	10.25	Apr. 23	<u>e/</u> 11.20	Sept. 2	12.68
Oct. 30	10.12	Apr. 29	11.05	Sept. 23	12.21
Nov. 5	10.44	May 7	11.92		
Dec. 3	11.51	May 14	12.20		
Dec. 10	11.53				
Jan. 10, 1933	11.48				

34. Mrs. W. E. Gould, $1\frac{1}{2}$ miles north-northeast Balmorhea postoffice.
Measuring point, benchmark in 1 X 12-inch plank, at south side of hole in plank, at ground level.

Date	Depth to water (feet)	Date	Depth to water (feet)	Date	Depth to water (feet)
Nov. 3, 1930	16.52	Dec. 10, 1932	11.85	May 7, 1933	13.28
Mar. 8, 1931	<u>e/</u> 16.94	Jan. 10, 1933	12.88	May 14	13.85
Nov. 3	16.67	Jan. 14	13.00	May 20	13.98
Dec. 11	17.40	Jan. 21	13.11	May 28	14.18
Jan. 6, 1932	<u>a/</u> 17.43	Jan. 28	13.22	June 4	14.35
Jan. 19	17.62	Feb. 4	13.24	June 11	14.42
Feb. 15	<u>b/</u> 17.00	Feb. 11	15.52	June 18	14.53
Apr. 13	16.63	Feb. 18	15.40	June 25	14.60
May 30	16.3	Feb. 25	<u>d/</u> 13.48	July 8	15.18
Aug. 4	17.05	Mar. 4	13.30	July 15	15.35
Sept. 17	<u>c/</u> 11.80	Mar. 11	13.51	July 29	15.70
Oct. 15	8.31	Apr. 1	13.42	Aug. 5	15.81
Oct. 22	9.05	Apr. 9	13.48	Aug. 11	16.26
Oct. 30	9.74	Apr. 16	13.60	Aug. 19	15.78
Nov. 5	10.22	Apr. 23	12.62	Aug. 26	15.62
Dec. 3	11.68	Apr. 29	12.90	Sept. 23	<u>g/</u> 14.65

a/ Rainfall from August 31 to January 1932, inclusive, was about 27 percent below 13-year average.

b/ Rainfall was 3.79 inches in February 1932, about 5 times the 13-year average.

c/ Rainfall was 15.24 inches in August and September 1932, more than 3 times the 13-year average.

d/ Rainfall in January and February 1933 was only .69 inch.

e/ Adjoining field under irrigation.

f/ Adjoining field recently irrigated.

g/ Rainfall April to August 1933, inclusive, was about one-third of 13-year average.

h/ Water running over top of well.

INTAKE OF GROUND WATER

Areas of intake

As has been pointed out in the section on the geology of this district (pp.10-12), the principal area of intake or replenishment for the large springs at Balmorhea is in a long narrow anticlinal valley which parallels the eastern escarpment of the Davis Mountains, west, northwest and southwest of the springs. In this valley the beveled edges of the Lower Cretaceous limestones appear at the surface or lie beneath a mantle of stream and terrace gravels not far below the surface and their honeycombed and cavernous members absorb and store a large part of the local rainfall and mountain run-off. The length of this intake area is not exactly known. It is believed, however, that the limestones may take in water along the anticline all the way from Big Aguja Canyon northwestward to the vicinity of San Martine in Pecos County, a distance of about 35 miles. (fig.1), p. 82.

The creeks from Aguja, Little Aguja, Madera and Cherry Canyons lose heavily between the mouths of their canyons and the downstream boundary off the anticline. The Lower Cretaceous rocks take in considerable water also in an area of outcrop along Limpia Creek about 20 miles southeast of Balmorhea. Practically all the discharge of Limpia Creek disappears during moderate and low stages in that locality.

The lavas of the Davis Mountains have an important part in the intake system. They absorb much of the mountain rainfall and run-off. The water moves downward through joints and crevices in the rocks until it reaches relatively impermeable layers of volcanic rock or tuff or underlying Upper Cretaceous clays. It then moves laterally and appears as springs or seeps in the mountain canyons and on the slopes of the eastern escarpment, where it is largely dissipated by evaporation and transpiration. A part, however, reaches the canyon streams, in places maintaining a small perennial flow, or is absorbed by the gravels of the streams and adjacent terraces. In at least one

locality considerable stream water apparently is absorbed by the volcanic rocks where synclinal structure has brought them below stream level. This is on Limpia Creek near old Limpia Post Office, about 15 miles south of Balmorhea. The water that enters there may rise into the stream gravels below the lower limb of the syncline and through them eventually reach the intake area of the lower Cretaceous rocks, 15 to 20 miles below.

The gravels also have an important part in the intake system. A large part of the district between Balmorhea and the mountains is underlain by stream and terrace gravels and in places tongues of these gravels extend a considerable distance into the mountains. The gravels serve as both a reservoir and a conduit. They absorb a part of the water that is discharged by the mountain seeps and springs and a part of the rainfall and run-off on the escarpment of the mountain front and the belt between the escarpment and outcrop of the Lower Cretaceous rocks. This water moves slowly in the gravels in a downstream direction until it reaches the outcrop of the Lower Cretaceous rocks and is gradually paid out to the reservoir in these rocks. The igneous rocks and gravels therefore, comprise an important part of the extensive storage and regulatory system by which the flow of the large springs is maintained during periods of months and even years of drought.

The water that supplies the flow of the smaller springs at Balmorhea is derived from the Toyah Valley gravels. It is replenished by rainfall on the valley floor and by seepage from Toyah Creek and from canals, ditches and irrigated lands and to some extent by seepage from the large springs.

Intake from Streams

Two gaging stations have been maintained on Limpia Creek, the upper station is Limpia Canyon near the site of the old Limpia Post Office about 16 miles southwest of Balmorhea and the other about 30 miles downstream on the bridge at the crossing of State Highway 190 (Old Spanish Trail) where the stream is commonly known as Barilla Creek or Draw.

The upper station was established February 27, 1925, and maintained until August 30, 1932, when it was destroyed by a flood. The lower station was installed December 6, 1924 and discontinued July 15, 1926. It was reestablished September 20, 1932, and a few days later, on September 29, was destroyed when the highway bridge was carried away by one of the largest floods that has occurred on this Creek in the memory of residents in this part of Texas. Neither station was rebuilt. Funds were not available to meet the high cost of construction of the type of gaging station necessary to withstand large floods.

Moreover, in the stretch in which the lower station is located, the shifting of the channel with each flood prevents the compilation of a satisfactory rating curve. The site of the upper station is above any known outcrop of the Lower Cretaceous limestones, while the lower site is below a section in which the streams cross several miles of such outcrop.

In 1932 and 1933 a program of stream measurements was undertaken on several of the mountain streams, including miscellaneous measurements on Limpia Creek, to obtain information both as to the volume of mountain run-off and the extent of seepage losses from the mountain streams in crossing the intake area of the Lower Cretaceous limestones. A permanent gaging station was established in Madera Canyon and equipped with a continuous water-stage recorder. Temporary staff gages were installed on the streams that issue from Cherry, Aguja and Little Aguja Canyons and read daily or weekly for several months. Numerous miscellaneous measurements of the discharge of these streams and Limpia Creek were made with a current meter and several series of such measurements were undertaken

on each stream to determine seepage losses on the intake area of the Lower Cretaceous limestones and gains and losses in various other sections, mostly below the mouths of the canyons. The most of the "seepage measurements" were made on a declining stage and therefore tend to show losses in each section that are smaller than the true losses or even to show a gain when actually there may have been a substantial loss. However, in making the measurements the stream gager moved in a downstream direction at a rate not much faster than the movement of the stream; in this way a large part of the error, due to changes in stage while the seepage measurements are progressing, can be eliminated and reasonably accurate estimates can be made of the losses and gains in the sections during comparable rates of stream discharge.

The records obtained at the gaging stations on Limpia and Barilla Creeks and in Madera Canyon have been published in various Water-Supply Papers of the Geological Survey. Most of the results of the measurements in 1932 and 1933 are given in Water-Supply Paper: 748, pages 140 to 160 and 171 to 174, 1933. These pages have been incorporated into this report (see pp. 56-80). The results of the measurements are discussed below by streams.

Madera Canyon.-- The gaging station in Madera Canyon is located about three-quarters of a mile above its mouth. The station was installed July 28, 1932, and has been in continuous operation ever since. According to the Fort Davis and Valentene topographic maps, the canyon drains about 54 square miles above the gage, consisting of mountain slopes and summits, mostly more than 5,500 feet above sea level. The daily records at the station have been published by the Geological Survey.^{1/} These records are fairly accurate for low and moderate

^{1/} Geological Survey Water-Supply Papers 748, 763, 788 and 808.

stages of stream discharge but comparatively poor for high stages. The estimated total annual discharge from 1932 to 1936 is given in the following table:

Discharge of Madera Canyon from
1932 to 1936, in acre-feet

Year (Oct. 1 - Sept. 30)	Discharge (acre-feet)
1932 (Aug. 1 to Sept. 30)	14,800
1932-33	6,000
1933-34	244
1934-35	637
1935-36	4,200

According to the daily record, the discharge from August to October 1932 amounted to about 17,000 acre-feet, (see table, p. 56) or approximately two-thirds of the total recorded discharge from 1932 to 1936. Altogether in the period of 4 years and 2 months there were 64 days in which the average discharge during 24 hours amounted to more than 25 second-feet, of which 25 occurred in August, September and October 1932, 18 in August and September 1933, one in July 1935 and 20 in May and September 1936. Most of the time the stream was dry or the discharge was 2 or 3 second-feet or less. The maximum discharge, somewhat more than 2,000 acre-feet, occurred on September 29 and 30, 1932.

Nine series of discharge measurements were made to determine seepage losses from Madera Canyon between the gage and points two to 10 miles below it. The maximum flow at the gage during these studies was 67.5 second-feet, the minimum 5.2 second feet and the average about 27 second-feet (table, p. 57). None of the water reached Toyah Creek, about 12 miles below the gage, and most of it was lost within a few miles. For example: On September 2, 1932, all but 2.2 second-feet of a discharge of 67.5 second-feet was lost within 8.2 miles of the gage, and on October 6, 1932 a discharge of 37.3 second-feet disappeared entirely within 4 miles. On August 27, 1933, a loss of only 6.1 second-feet was indicated in a stretch of 6.2 miles below the gage but these measurements were made

during a rapidly declining stage and the true loss probably was much greater. No determinations of seepage losses were made during flood stages when the discharge amounted to several hundred second-foot, but at such times, the water spreads over a wide expanse of gravels and the rate of loss must be heavy.

Cherry Canyon.- Cherry Canyon drains about 70 square miles of mountainous area as compared with about 55 miles by Madera Canyon, and the drainage area is of similar ruggedness and altitude. A Staff gage was installed on the stream about half a mile below the mouth of the Canyon on August 1, 1932, and read daily until November 13, and again from August 30 to September 9, 1933. In the intervening period, readings were made once a week. From this fragmentary record of gage heights, supplemented by miscellaneous measurements with a current meter, the conclusion is reached that the discharge of the Canyon from August 1, 1932 to September 30, 1933 was materially greater than the discharge of Madera Canyon during corresponding periods. The seepage studies disclosed that the stream disappears within a relatively short stretch below the gage during low and moderate stages. For example: On September 15, 1932 a discharge of 13.5 second-feet, and on October 7, 1932, a discharge of 31.2 second-feet was entirely lost within 2-1/2 miles of the gage. (table, p. 74a)

Big Aguja Canyon. - Numerous measurements were made in 1932 and 1933, of the discharge of this Canyon, at or near the dam of the Texas Pacific Railway; the largest discharge, computed at more than 4,000 second-feet, being on September 7, 1932. (table, pp.77-78).

A staff gage was installed in the Canyon on July 27, 1932 and read intermittently until September 1, 1933. The area drained above the gage consists of about 47 square miles, as compared with 54 square miles above the gaging station in Madera Canyon. The two areas apparently are similar

topographically. The records indicate that the discharge of the Canyon was greater than the discharge from Madera Canyon in the early part of September 1932, and somewhat less in the latter part of that month, but the record in Big Aguja Canyon is too fragmentary and too short to justify any conclusions as to whether the run-off over a long period of time is comparable.

Seepage surveys were made on September 1, 3 and 13, and October 6, 1932 (table, p. 60) to determine losses between the gage and Toyah Creek, 9.6 miles below the gage. On September 1, the entire flow amounting to 18.7 second-feet was lost before Toyah Creek was reached. Similarly, on September 13, the total flow of 8.6 second-feet was lost. On October 6, the loss was 18.4 second-feet including the discharge at the gage plus inflow from tributaries below. On September 3, only 26.6 second-feet was lost from a discharge of 45.4 second-feet, but the measurements on that date were taken on a rapidly declining stage and the actual loss probably was greater.

Little Aguja Canyon.- Numerous measurements of the discharge of this canyon and its South Fork were made in 1932 and 1933 (table, p. 77), the maximum discharge from the main canyon being estimated as 2,640 second-feet on August 29, 1932. The stage of the stream was recorded daily for a few weeks in the latter part of the summer in 1932 at a temporary staff gage, at a point in the canyon above which approximately 31 square miles are drained. From the gage height record and a few current meter measurements, it is estimated that the discharge of the canyon during July and August may have been about 8,000 acre-feet or a little more than half the discharge of Madera Canyon.

Seepage surveys were made on this stream August 17, September 1, 13, 14, and 20, and October 6, 1932. In these surveys, it was found that the

entire discharge of the stream ranging from 1.3 to 26.8 second-feet was lost before Toyah Creek was reached. (table p. 59).

Limpia Creek.-- The yearly discharge of Limpia Creek at the gaging station in Limpia Canyon is given below:

Discharge of Limpia Creek in Limpia Canyon
16 miles southwest of Balmorhea
Drainage area - 272 square miles

Year (October 1 to September 30)	Discharge (acre-feet)
1925 (Feb. 27 - Sept. 30)	4,080
1925-26	577
1926-27	661
1927-28	7,160
1928-29	839
1929-30	894
1930-31	1,610
1931-32 (October 1 to Aug. 27)	2,590

The station was destroyed by a flood August 30, 1932, in which it is estimated a peak discharge of about 14,000 second-feet was reached. The stream was measured at the lower gaging station from December 6, 1924 to July 15, 1926, where as explained on p. 43, it is commonly known as Barillo Creek or Draw. During most of that period, there was no discharge at either station, but from March to September 1925, several runs of storm water occurred in which the discharge amounted to a total of about 4,800 acre-feet at the upper station and about 11,000 acre-feet at the lower one. The sites of the two stations are about 40 miles apart and the drainage from about 100 square miles of mountains, mostly less than 5,000 feet in altitude, enters the stream between them.

Five series of current meter measurements were made on Limpia Creek to determine seepage losses and gains in various sections above and below the site

of the gaging station in Limpia Canyon. These measurements indicate that the stream increases from seepage inflow from the vicinity of Fort Davis down to a point about 2 miles above the gaging site. Below that point, the stream loses in several sections and disappears entirely during low and moderate stages before reaching observation points, 19 to 22 miles below the station, and 9 to 12 miles below the Jeff Ranch. (fig. 1) The loss in the vicinity of the gaging station was as follows:

October 8, 4.2 second-feet; November 1, 8.6 second-feet; November 21, 4.1 second-feet; August 3, 1933, 12.6 second-feet. The loss between the Jeff Ranch and points of measurement, a half mile to one mile below, was 6.2 feet on November 1, 1932 and 2.4 second-feet on November 21, 1932. On October 8, 1932, a flow of 65.1 second-feet disappeared between the Jeff Ranch and the observation point, 12 miles below.

Discharge of ground water.

Spring discharge.- The table below, compiled from the gaging records, gives the approximate mean discharge of Phantom Lake, San Solomon, Giffin, West Sandia and East Sandia Springs for the following periods: (1) Start of record in the fall of 1931 to August 1932, inclusive; (2) September to December 1932, inclusive, and (3) January 1933 to end of record in September 1933.

Mean discharge of Phantom Lake, San Solomon, Giffin, ^{1/}
West Sandia and East Sandia Springs, in second-feet, 1931-33

	1 Oct., Dec. 1931 to Aug. 1932	2 Sept. to Dec. 1932	3 Jan. 1933
Phantom Lake	13	35	18
San Solomon	33	60	40
Giffin	4.7	5.6	5.0
West Sandia	1.1	1.7	1.3
East Sandia	1.1	13	1.2
Total	53	115	65

^{1/} See pages 61 to 67 and 72 to 74 for records of daily discharge.

The mean discharge of the five springs amounted to 53 second-feet, or about 23,800 gallons a minute, in the first period, 104 second-feet, or about 46,700 gallons a minute in the second; and 65 second-feet, or about 29,300 gallons a minute in the third. In the first period the rainfall at Balmorhea was materially above the average, due to a fall of 2.25 inches in December 1931 and 3.79 inches in February 1932. In the second period, the rainfall was abnormally high, amounting to 11.64 inches in September 1932 alone, or about 85 percent of the average annual rainfall. In the third period, the rainfall was about 40 percent below the average. The total discharge during the three periods, aggregating almost 2 years, was at the average rate of about 48,000 acre-feet a year.

The records of the combined discharge of the Saragosa and Toyah Creek springs are too few to warrant making an estimate of their mean discharge during the same periods, but the available measurements indicate that it may have been around 9 to 10 second-feet or at the rate of 6,500 to 7,000 acre-feet a year. Thus the flow of all the springs from the fall of 1931 to the fall of 1933 probably averaged about 54,000 to 55,000 acre-feet a year.

Since approximately 25,000 acre-feet of the discharge occurred in the four months September to December 1932, following abnormally heavy rains, the mean discharge of the springs, during the period of record, must have been materially greater than their long time average discharge.

Analyses of water from gravity springs near Balmorhea, Texas

(Parts per million.)

Analyzed by E. W. Lohr and L. A. Shinn

Spring	Date of collection	Total dissolved solids	Calcium (Ca)	Magnesium (Mg)	Sodium and Potassium (Na+K) (calc.)	Bicarbonate (HCO ₃)	Sulfate (SO ₄)	Chloride (Cl)	Nitrate (NO ₃)	Total hardness as CaCO ₃
Toyah Creek	Nov. 13, 1931	--	200	81	463	222	705	655	--	832
Saragosa	Dec. 7, 1930	2,846	272	102	584	332	868	842	5.0	1,098
East Sandia	Dec. 7, 1930	2,999	276	101	631	309	992	840	6.0	1,104

Partial analyses of water from wells in vicinity of Balmorhea, Texas

(Parts per million.)

(Well numbers correspond with numbers in table of well records.)

(Analyzed by E. W. Lohr and L.A. Shinn)

Well No.	Owner	Depth of well (ft.)	Date of collection	Total dissolved solids (calc.)	Calcium (Ca)	Magnesium (Mg)	Sodium and Potassium (Na+K) (calc.)	Bicarbonate (HCO ₃)	Sulfate (SO ₄)	Chloride (Cl)	Nitrate (NO ₃)	Total hardness as CaCO ₃ (calc.)
1	Balmorhea Livestock Company	16	Dec. 7, 1930	2,770	325	67	560	334	849	810	5.0	1,087
2	A. C. Schreyer	14	Dec. 7, 1930	2,293	230	73	471	288	706	655	5.0	874
4	A. W. Wigley	20	Nov. 13, 1931	--	370	145	787	438	1,174	1,170	--	1,520
6	Sedro Garcia	20	Nov. 13, 1931	--	306	86	574	330	881	835	--	1,118
11	Mrs. Nell B. Westerman	20	Nov. --, 1931	--	332	126	764	324	1,131	1,110	--	1,347
12	J. P. Hill	22	Nov. 13, 1931	--	300	143	721	302	1,366	875	--	1,336
13	Rosenbaum Company	15	Nov. 13, 1931	--	305	109	699	349	1,046	960	--	1,209
17	Do.	11	Nov. 13, 1931	--	322	93	590	290	965	870	--	1,186
18	Mrs. Paul Renz	12	Nov. 13, 1931	--	347	91	551	268	858	940	--	1,241
24	E. P. Stuckler	13	Nov. 13, 1931	--	319	144	849	382	1,291	1,118	--	1,388
27	B. W. Vanderson	20	Nov. 13, 1931	--	273	114	647	293	1,031	882	--	1,150
28	E. W. Backus	18	Nov. 13, 1931	---	350	155	792	320	1,491	1,005	--	1,510
30	W. E. Gould	36	Dec. 7, 1930	1,342	301	35	111	337	363	335	12	896
37	Saragosa School	158	Dec. 6, 1930	3,283	346	137	595	334	1,009	982	15	1,427

Conclusions

The Phantom Lake, San Solomon and Giffin Springs come from an extensive system of solution channels in lower Cretaceous limestone, comprising in the aggregate a very large reservoir. The Lower Cretaceous rocks are underlain by rocks of Permian age, which yield to wells in other parts of the basin of Toyah Creek water that is rather highly mineralized. Water enters the Lower Cretaceous limestone along the front of the Davis Mountains and probably all or nearly all of the water is discharged by the springs. The limestone is close to the surface along the northeast edge of a syncline at the Phantom Lake Spring. A few hundred feet east of the spring relatively impermeable strata of upper Cretaceous age are dropped down against the limestone on the northeast side of a fault, thereby creating a barrier which causes the water to rise to the surface in the spring. The throw of this fault is not very great and a large part of the underground water escapes over the crest of the barrier and continues underground toward the San Solomon and Giffin springs. In the vicinity of the San Solomon spring the limestone again comes nearly to the surface and a short distance to the east of the springs the Upper Cretaceous strata appear in such a way that it seems necessary to assume that they are again dropped down against the limestone on the northeast side of a fault. This fault presumably is of considerably greater throw than the one immediately below the Phantom Lake Spring. It may create a practically complete barrier to the movement of the water in the limestone and bring all or nearly all of the flow to the surface in the San Solomon and Giffin springs.

The Saragosa, West Sandia, East Sandia, and Toyah Creek springs have their source in the relatively thin surficial gravel. In the valley between Toyahvale and the Brogado Hills the maximum depth of the gravels is likely to be 60 feet or less, and in most places it is less than 40 feet. Between the Brogado Hills and Saragosa the gravel and associated sand, silt, and clay have a thickness of several hundred feet; there the deposits are not very permeable and the water table is low.

A well put down in the vicinity of Phantom Lake or San Solomon Springs above the faults might tap the limestone reservoir at shallow depths, but it would be practically certain to decrease the flow of the springs. Between the San Solomon Spring - Giffin Springs fault and the Brogado Hills the water-bearing horizon in the Lower Cretaceous limestone probably is from 500 to 1,000 feet beneath the surface, perhaps around 700 or 800 feet. If the fault barrier just below the springs is very tight the withdrawal of water from the limestone in this stretch by means of a well would not affect the flow of the springs. If the barrier is not complete, a well in that area would tend to decrease the flow of the springs. In this connection, the following possibility should be kept in mind: If the fault below the springs offers an effective barrier to the down-valley movement of the water underground, solution channelling may not have developed extensively in the limestone below the barrier because of lack of circulation. As a result, the limestone may be relatively tight between the San Solomon Spring and the Brogado Hills and the water in it may be comparatively highly mineralized.

Before a well is put down in this part of the valley the San Solomon and Givvin Springs should be equipped with automatic measuring devices and accurate continuous records kept of their flow before the well is placed in operation and while it is being operated to determine whether the well affects the springs.

Shallow wells in the gravels in parts of the Balmorhea district may yield a few second-feet of water but such wells, if improperly located, or pumped too heavily may deplete the flow of the Toyah Creek, Saragosa or Sandia Springs. The flow of these springs is effectively utilized only during the growing season and in view of this and of the urgent need for an additional supply of water during the relatively short critical period of the growing season, an attempt to develop shallow ground water would appear to be justified. If the wells were put down near the upper end of the district three to four miles southwest of Balmorhea, it is possible that the withdrawal of water from them would not seriously affect the flow of the springs until after the critical period of the growing season. Then during the succeeding months the normal recharge to the shallow ground-water reservoir would be expected to replenish the supply in storage in the gravel. Some water might also be developed by shallow wells in the moist area, subject to seepage from the Balmorhea irrigation reservoir, between the reservoir and Balmorhea. If wells are sunk in the gravel and pumped for irrigation, especially in the district near Balmorhea, the development should proceed cautiously and in the meantime continuous accurate records should be kept of the flow of the Saragossa, Sandia, and Toyah Creek springs, and of the position of the water table.

In the district between the Brogado Hills and Saragosa the limestone is at greater depth, perhaps around 1,000 to 1,200 feet. It is reasonably probable that any water that occurs in the limestone in that area is shut off from the reservoir that supplies the large springs at Balmorhea, and has its source partly in separate areas of intake and partly in the overflow that passes from those springs. A deep well in this district, therefore, would not be expected to interfere with the springs.

A deep well in the vicinity of Balmorhea, perhaps 1,200 to 1,300 feet, would probably reach a water-bearing horizon in the basal Lower Cretaceous sands or in the Permian rocks. The quality of the water that might be encountered in these rocks is, however, not known.

Records of the discharge of the streams and springs and seepage gains and losses from the streams and canals of the Balmorhea district are given in the following pages. These records are taken from Water-Supply Paper 748.

Miscellaneous Measurements

of the

Discharge of San Solomon, Giffin, Saragosa and Toyah Creek springs
Near Balmorhea, Texas in second-feet

Date	San Solomon Spring	Giffin Spring	Saragosa and Toyah Creek Springs
Aug. 20, 1919	34.7		4.8
July 20, 1922	37.0	3.9	5.9
April 26, 1923	26.5	5.4	
Sept. 25, 1923	35.0		
June 3, 1924	32.3	3.9	
June 4, 1924	33.4		
Sept. 6, 1924	34.4		
Sept. 7, 1924	35.4		
Oct. 17, 1924	37.7		
Jan. 8, 1925	38.2		
Mar. 4, 1925	36.3	2.9	
June 2, 1925	34.3	3.4	
Feb. 6, 1935	30.4		
July 2, 1935	32.8		
Aug. 13, 1935	31.6		
April 15, 1936	30.4		

Discharge, in second-feet, of Creek in
Madera Canyon near Toyahvale, Tex.

1932-33

Day	Aug.	Sept.	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	June	July	Aug.	Sept.
1	0	21	530	2.1	1.4	1.1	0.2	0.5	0	3.5	0	102
2	0	54	141	2.1		.9	.2	.5	0	2.6	0	119
3	0	85	108	2.1		.8	.2	.4	0	2.0	0	44
4	0	38	51	2.1		.7	.2	.4	0	.7	0	24
5	0	42	40	2.1		.7	.2	.4	0	.2	0	19
6	0	87	40	2.1		.7	.2	.4	0	.1	0	13
7	0	659	50	2.1		.7	.2	.3	0	.1	0	9.1
8	0	351	22	1.4		.6	.2	.3	0	0	0	7.2
9	0	138	16			.6	.2	.3	0	0	0	6.9
10	0	138	11			.6	.2	.3	0	0	0	103
11	0	54	16			.5	.2	.2	0	0	0	104
12	0	30	16			.5	.2	.1	0	0	0	67
13	0	14	11			.5	.2	.1	0	0	0	58
14	0		7.5			.4	.2	.1	0	0	0	70
15	0	14	7.5		1.2	.4	.2	.1	0	0	0	39
16	0	14	16		1.1	.4	.2	.1	0	0	0	27
17	0	14	7.5		1.1	.4	.2	.1	0	0	0	19
18	0	14	7.5		1.1	.4	.2	.1	0	0	0	15
19	0	9.0	7.5		1.4	.9	.4	.2	.1	0	0	0
20	0	5.1	7.5	.7		.5	.2	.1	0	0	0	139
21	0	5.1	7.5	.7		.2	.2	.1	0	0	0	28
22	0	5.1	7.5	.8		.2	.2	0	0	0	0	16
23	0	9.0	7.5	2.3		.2	.1	0	0	0	0	11
24	0	9.0	7.5	2.1		.2	.1	0	0	0	5.3	7.9
25	0	9.0	7.5	1.8		.3	.3	0	0	0	24	5.6
26	0	9.0	7.5	1.6		.3	.4	0	0	0	11	4.4
27	0	9.0	7.5	1.3		.3	.5	0	0	0	124	3.5
28	27	134	4.4	1.3		.3	.5	0	17	0	78	3.0
29	321	2,190	4.4	1.2		.3		0	5.9	0	42	2.6
30	810	2,040	4.4	1.6		.3		0	5.8	0	213	2.1
31	94		4.4		1.3	.5		0		0	60	

Month	Maximum	Minimum	Mean	Run-off in acre-feet
1932				
July 28-31	0	0	0	0
August	810	0	40.4	2,480
September	2,190	5.1	206	12,500
The period				14,800
1932-33				
October	530	4.4	37.5	2,310
November			1.56	93
December			1.35	83
January	1.1	.2	.47	29
February	.5	.1	.22	12
March	.5	0	.16	9.8
June	17	0	.96	57
July	3.5	0	.30	18
August	213	0	18.0	1,110
September	139	2.1	38.3	2,280
The year	530	0	8.28	6,000

Note. - No flow during April and May 1933.

Discharge measurements in Madera Canyon to determine losses from
seepage from a point 13.3 miles above to a point 3.5 miles above
Toyahvale, Tex., 1932-33

Date	Stream	Location	Distance from initial point (miles)	Discharge (second-feet)			
				Main stream	Tribu- tary	Gain or loss in section	Total gain or loss
1932							
Sept. 1	Madera Canyon	Gage	13.3	23.2	-	-	-
1	do.	Madera Springs road crossing	7.1	3.9	-	-19.3	-19.3
1	do.	8.2 miles below gage	5.1	0	-	- 3.9	-23.2
2	do.	Gage	13.3	67.5	-	-	-
2	do.	Madera Springs road crossing	7.1	41.8	-	-25.7	-25.7
2	do.	8.2 miles below gage	5.1	2.2	-	-39.6	-65.3
2	do.	Duncan Kingston crossing	3.5	.5	-	- 1.7	-67.0
3	do.	Gage	13.3	41.8	-	-	-
3	do.	Madera Springs road crossing	7.1	16.1	-	-25.7	-25.7
3	do.	8.2 miles below gage	5.1	8.3	-	- 7.8	-33.5
3	do.	Duncan Kingston crossing	3.5	4.5	-	- 3.8	-37.3
12	do.	Gage	13.3	28.7	-	-	-
12	Madera Springs Creek	Mouth	11.3	-	0.5	-	-
12	Madera Canyon	Madera Springs road crossing	7.1	7.2	-	-22.0	-22.0
12	do.	8.2 miles below gage	5.1	2.5	-	- 4.7	-26.7
15	do.	Gage	13.3	14.5	-	-	-
15	do.	Madera Springs road crossing	7.1	.3	-	-14.2	-14.2
15	do.	8.2 miles below gage	5.1	.2	-	- .1	-14.3
21	do.	Gage	13.3	5.2	-	-	-
21	do.	Rock outcrop	7.5	0	-	- 5.2	- 5.2
21	do.	Madera Springs road crossing	7.1	.3	-	✓ .3	- 4.9
21	do.	0.2 miles below Madera Springs road crossing	6.9	0	-	- .3	- 5.2
21	do.	8.2 miles below gage	5.1	.6	-	✓ .6	- 4.6
24	do.	Gage	13.3	6.7	-	-	-
24	Side canyon	Mouth	13.2	-	.3	-	-
24	Madera Canyon	1.5 miles below gage	11.8	6.7	-	- .3	- .3
24	do.	2.5 miles below gage	10.8	1.0	-	- 5.7	- 6.0
24	do.	3.5 miles below gage	9.8	.2	-	- .8	- 6.8
Oct. 6	do.	Gage	13.3	37.3	-	-	-
6	Madera Springs Creek	Mouth	11.3	-	1.5	-	-
6	Madera Canyon	2.6 miles below gage	10.7	22.6	-	-16.2	-16.2
6	do.	3.7 miles below gage	9.6	4.5	-	-18.1	-34.3
6	do.	Limestone outcrop	9.3	0	-	- 4.5	-38.8

Discharge measurements in Madera Canyon--Continued
1932-33

Date	Stream	Location	Distance from initial point (miles)	Discharge (second-feet)			
				Main stream	Tribu- tary	Gain or loss in section	Total gain or loss
1933							
Aug. 27	Madera Canyon	Gage	13.3	17.5	-	-	-
27	do.	Madera Springs road crossing	7.1	11.4	-	- 6.1	- 6.1
27	do.	Duncan Kingston crossing	3.5	0	-	-11.4	-17.5

Discharge measurements of Toyah Creek to determine losses from
seepage from a point 1.2 miles above to a point 8.8 miles below
Toyahvale, Tex., 1932-33

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)				
				Main stream	Tribu- tary	Diver- sion	Gain or loss in section	Total gain or loss
Nov. 6	Toyah Creek	Aloma settlement	1.0	13.6	-	-	-	-
6	Project waste	0.9 mile above Balmorhea	3.9	-	0.3	-	-	-
6	Saragosa Springs Creek	150 feet above mouth	4.7	-	9.9	-	-	-
6	Toyah Creek	500 feet below Balmorhea Bridge	4.8	29.1	-	-	+5.3	+5.3
6	do.	500 feet below Moore Dam	5.8	30.7	-	-	+1.6	+6.9
6	do.	Saragosa Dam	8.8	29.0	-	-	-1.7	+5.2
Jan. 23	do.	1.8 miles above Balmorhea Bridge	3.0	0	-	-	-	-
23	Saragosa Springs Creek	200 feet above mouth	4.7	-	8.2	-	-	-
23	Toyah Creek	300 feet below Balmorhea Bridge	4.8	12.5	-	-	+4.3	+4.3
23	do.	300 feet below Moore Dam	5.8	12.8	-	-	+ .3	+4.6
23	do.	Saragosa Dam	8.8	10.5	-	-	-2.3	+2.3
Mar. 14	do.	United States Highway 290 crossing	-1.2	2.5	-	-	-	-
14	do.	1.8 miles above Balmorhea	3.0	0	-	-	-2.5	-2.5
14	Saragosa Springs Creek	200 feet above mouth	4.7	-	6.7	-	-	-
14	Toyah Creek	500 feet below Balmorhea Bridge	4.8	9.4	-	-	+2.7	+ .2
14	Moore Canal	150 feet below takeout	5.8	-	-	5.3	-	-
14	Toyah Creek	500 feet below Moore Dam	5.8	4.8	-	-	+ .7	+ .9
14	Saragosa Canal	50 feet below takeout	8.8	-	-	6.0	-	-
14	Toyah Creek	50 feet below Saragosa Dam	8.8	.2	-	-	+1.4	+2.3
May 16	do.	1.8 miles above Balmorhea Bridge	3.0	0	-	-	-	-
16	Saragosa Springs Creek	50 feet above mouth	4.7	-	6.4	-	-	-
16	Toyah Creek	500 feet below Balmorhea Bridge	4.8	9.2	-	-	+2.8	+2.8
16	Moore Canal	2,000 feet below Moore Dam	5.8	-	-	7.3	-	-
16	Toyah Creek	500 feet below Moore Dam	5.8	1.6	-	-	- .3	+2.5
16	Saragosa Canal	50 feet below Saragosa Dam	8.8	-	-	4.0	-	-
16	Toyah Creek	do.	8.8	0	-	-	+2.4	+4.9
July 11	do.	1.8 miles above Balmorhea Bridge	3.0	0	-	-	-	-
11	Saragosa Springs Creek	200 feet above mouth	4.7	-	5.6	-	-	-
11	Toyah Creek	500 feet below Balmorhea Bridge	4.8	8.3	-	-	+2.7	+2.7
11	Moore Canal	2,000 feet below Moore Dam	5.8	-	-	7.0	-	-
11	Toyah Creek	500 feet below Moore Dam	5.8	1.6	-	-	+ .3	+3.0
11	Saragosa Canal	125 feet below Saragosa Dam	8.8	-	-	3.5	-	-
11	Toyah Creek	50 feet below Saragosa Dam	8.8	0	-	-	+1.9	+4.9

Discharge measurements in Little Aguja Canyon to determine losses from
seepage from a point 15.5 miles above to a point 2.2 miles above
Toyahvale, Tex., 1932

Date	Stream	Location	Distance from initial point (miles)	Discharge (second-feet)			
				Main stream	Tribu- tary	Gain or loss in section	Total gain or loss
Aug. 17	Little Aguja Canyon	Temporary staff gage	15.5	1.3	-	-	-
17	do.	2.8 miles below staff gage	12.7	1.0	-	-0.3	-0.3
17	South Fork of Little Aguja Canyon	0.2 mile above mouth	11.5	-	0.4	-	-
17	Little Aguja Canyon	5.5 miles below gage	10.0	.4	-	-1.0	-1.3
17	do.	7.0 miles below gage	8.5	.6	-	/.2	-1.1
17	do.	8.5 miles below gage	7.0	.4	-	-.2	-1.3
17	do.	9.6 miles below gage	5.9	.2	-	-.2	-1.5
17	do.	300 feet above limestone bluff	4.3	.1	-	-.1	-1.6
17	do.	Upper end of limestone bluff	4.3	0	-	-.1	-1.7
17	do.	Lower end of limestone bluff	4.0	.2	-	/.2	-1.5
17	do.	Mouth	2.2	0	-	-.2	-1.7
Sept. 1	do.	3 miles below gage	12.5	12.1	-	-	-
1	South Fork of Little Aguja Canyon	0.2 mile above mouth	11.5	-	6.4	-	-
1	Little Aguja Canyon	Mouth	2.2	0	-	-18.5	-18.5
13	do.	50 feet below mouth of South Fork	11.3	22.8	-	-	-
13	do.	100 feet above limestone bluff	4.3	0	-	-22.8	-22.8
13	do.	50 feet above limestone bluff	4.3	.1	-	/.1	-22.7
13	do.	0.1 mile below limestone bluff	3.9	0	-	-.1	-22.8
14	do.	50 feet below mouth of South Fork	11.3	11.2	-	-	-
14	do.	0.5 mile below white bluff	8.2	0	-	-11.2	-11.2
20	do.	50 feet below mouth of South Fork	11.3	4.1	-	-	-
20	do.	150 feet above upper Duncan road	9.5	0	-	-4.1	-4.1
20	do.	100 feet above white bluff	8.7	.6	-	/.6	-3.5
20	do.	600 feet below white bluff	8.6	0	-	-.6	-4.1
20	do.	Second white bluff	7.8	.6	-	/.6	-3.5
20	do.	0.1 mile below second white bluff	7.7	0	-	-.6	-4.1
Oct. 6	do.	200 feet below mouth of South Fork	11.2	26.8	-	-	-
6	do.	Lower Duncan road crossing	6.0	26.5	-	-.3	-.3
6	Wet-weather springs	Limestone bluff	4.3	-	.2	-	-
6	Little Aguja Canyon	50 feet below limestone bluff	3.9	2.2	-	-24.5	-24.8
6	do.	4,000 feet above mouth	3.0	0	-	-2.2	-27.0

Discharge measurements in Big Aguja Canyon to determine losses from seepage from a point 11.8 miles above to a point 2.2 miles above Toyahvale, Tex., 1932

Date	Stream	Location	Distance from initial point (miles)	Discharge (second-feet)		
				Main stream	Tributary	Gain or loss in section Total gain or loss
Sept. 1	Big Aguja Canyon	Temporary staff gage	11.8	18.7	-	-
1	do.	Above mouth of Seven Springs Creek	7.3	4.4	-	-14.3
1	Seven Springs Creek	Mouth	7.2	-	0.5	-
1	Big Aguja Canyon	do.	2.2	0	-	-4.9
3	do.	Temporary staff gage	11.8	45.4	-	-
3	do.	Above mouth of Seven Springs Creek	7.3	35.7	-	-9.7
3	Seven Springs Creek	Mouth	7.2	-	.4	-
3	Big Aguja Canyon	do.	2.2	19.2	-	-16.9
13	do.	Temporary staff gage	11.8	8.6	-	-
13	Walnut Canyon	Mouth	7.5	-	.2	-
13	Big Aguja Canyon	Above mouth of Seven Springs Creek	7.3	5.6	-	-3.2
13	Seven Springs Creek	Mouth	7.2	-	4.8	-
13	Big Aguja Canyon	do.	2.2	0	-	-10.4
Oct. 6	do.	Temporary staff gage	11.8	10.3	-	-
6	do.	Above pipe-line crossing	9.6	5.6	-	-4.7
6	Break in Texas & Pacific pipe line	Canyon crossing	9.5	-	1.5	-
6	Big Aguja Canyon	Below pipe-line crossing	9.5	7.1	-	0
6	Walnut Canyon	Mouth	7.5	-	.6	-
6	Big Aguja Canyon	Above mouth of Seven Springs Creek	7.3	14.0	-	-6.3
6	Seven Springs Creek	Mouth	7.2	-	6.1	-
6	Big Aguja Canyon	3 miles above mouth	5.2	0	-	-20.1
6	do.	$\frac{1}{2}$ mile above mouth	2.7	0	-	0
6	do.	0.4 mile above mouth	2.6	.1	-	-
6	do.	Mouth	2.2	.1	-	0

Discharge, in second-feet, of Phantom Lake Spring
near Toyahvale, Tex.
1931-33

Day	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1931-32										
1	-	13	15	20	16			12	13	b 18
2	-	13	13	19	16		13	12	12	b 18
3	-	15	15	17	16		13	13	12	18
4	-	15	15	16	16		13	13	12	b 19
5	-	13	15	16	16		13	12	12	b 23
6	-	13	13	18	15		13	12	12	b 26
7	-	15	15	16	14		15	12	12	34
8	-	15	15	20	14	a 14	15	13	12	42
9	-	15	15	21	14		13	13	13	45
10	-	15	15	22	14		13	13	13	45
11	-	15	13	22	14		15	12	13	45
12	-	14	13	22	14		13	13	15	46
13	-	15	15	22	b 14		13	13	15	45
14	-	13	13	22			13	13	13	42
15	-	13	15	20			15	15	13	40
16	-	15	13	19			13	13	13	37
17	-	13	13	19			15	15	15	35
18	-	15	13	18			13	13	15	34
19	-	13	15	17			15	15	15	33
20	-	13	13	17	a 14		13	13	13	b 32
21	b 13	13	13	b 17			15	15	12	31
22		13	15	b 17		a 15	13	13	12	30
23		13	20	b 17			13	13	12	29
24	a 13	13	19	b 16			13	13	12	b 26
25		13	22	b 16			13	13	12	26
26		13	24	b 16			12	13	12	26
27		13	25	b 16			12	13	12	25
28		13	24	b 16			12	15	12	38
29	b 13	15	22	16			12	15	12	34
30	b 13	13	-	16			12	15	15	54
31	13	13	-	16			-	13	17	39

a Estimated.
b Partly estimated.

Discharge, in second-feet, of Phantom Lake Spring near Toyahvale, Tex., 1931-33-
Continued

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1932-33												
1	104	31	21	18	17	16	16	16	15	c 15	14	} a 42
2	114	30	21	18	17	16	16	16	16	c 15	14	
3	114	30	21	18	17	16	16	17	c 16	15	14	
4	109	29	21	18	17	16	16	17	c 16	15	14	47
5	102	28	21	18	17	16	16	17	c 16	15	14	46
6	88	28	21	18	17	16	16	17	c 16	15	14	46
7	82	27	21	18	17	16	16	c 17	c 16	c 15	14	44
8	75	26	21	18	17	16	16	c 17	16	c 15	14	43
9	73	26	21	18	17	16	16	c 17	16	c 15	14	43
10	70	26	20	18	17	16	16	c 17	16	15	14	42
11	67	25	20	18	17	16	16	17	16	14	14	42
12	64	25	20	18	17	16	16	17	15	14	14	43
13	63	25	20	18	17	16	16	17	15	14	14	43
14	61	24	20	18	17	16	16	16	15	14	c 14	-
15	60	24	20	18	17	16	16	16	15	14	14	-
16	57	24	19	18	17	16	16	16	15	14	c 14	} a 36
17	55	24	19	18	17	16	16	16	15	14	14	
18	53	23	19	18	17	16	16	16	15	14	14	
19	51	23	19	18	17	16	16	16	14	14	14	-
20	49	23	19	18	17	16	16	16	14	14	14	c 28
21	46	23	19	18	17	16	16	16	14	14	14	c 28
22	44	23	19	18	17	16	16	16	15	14	14	c 28
23	43	23	19	18	17	16	16	16	15	14	14	27
24	42	23	19	18	17	16	16	16	15	14	14	26
25	39	23	19	18	17	16	16	16	15	14	c 14	25
26	38	22	19	18	16	16	16	16	15	14	c 14	24
27	37	22	19	18	16	16	16	16	15	14	c 14	22
28	35	22	19	17	16	16	16	16	c 15	14	14	21
29	33	21	19	17	-	16	16	15	c 15	14	16	21
30	32	21	19	17	-	16	16	15	c 15	14	a 24	21
31	31	-	18	17	-	16	-	15	-	c 14	-	-

Day	Oct.	Nov.	Dec.	Day	Oct.	Nov.	Dec.	Day	Oct.	Nov.	Dec.
1933				1933				1933			
1	20	16	} a 16	11	17	16	} a 16	21	16	16	a 16
2	20	16		12	17	15		22	16	16	16
3	20	16		13	17	15		23	16	} a 16	16
4	19	16		14	17	15		24	16		16
5	19	16		15	17	16		25	16		16
6	19	16	} a 16	16	17	16	} a 16	26	16	} a 16	16
7	19	16		17	17	16		27	16		16
8	18	16		18	17	16		28	16		16
9	18	16		19	17	16		29	16		16
10	17	16		20	16	16		30	16		16

a Estimated

c Partly estimated or interpolated.

Discharge, in second-feet, of Phantom Lake Spring near Toyahvale, Tex., 1931-33-
Continued

Month	Maximum	Minimum	Mean	Run-off in acre-feet
1931-32				
December 21-31	-	-	13.0	284
January	14	13	13.0	799
February	25	13	15.3	880
March	22	16	18.3	1,130
April	-	-	14.4	857
May	-	-	13.5	830
June	13	12	12.8	762
July	15	12	12.8	787
August	17	12	12.7	781
September	39	18	34.3	2,040
The period				9,150

Discharge, in second-feet, of Phantom Lake Spring near Toyahvale, Tex., 1931-33-
Continued

Month	Maximum	Minimum	Mean	Run-off in acre feet
1932-33				
October	114	31	62.3	3,830
November	31	21	24.8	1,480
December	21	18	19.7	1,210
January	18	17	17.9	1,100
February	17	16	16.9	939
March	16	16	16.0	984
April	16	16	16.0	952
May	17	15	16.3	1,000
June	16	14	15.2	904
July	15	14	14.3	879
August	-	14	14.7	904
September	-	21	25.1	2,090
The year	114	14	22.5	16,300
1933				
October	20	16	17.2	1,060
November	-	-	15.9	946
December	-	-	16.0	984
The year				2,990

Discharge, in second-feet, of Giffen Springs
at Toyahvale, Tex.
1931-33

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1931-32												
1	-	4.3	a3.7	3.6	4.5	4.6	4.7	4.7	4.6	4.7	a4.7	a4.7
2	-	4.2	a3.7	3.6	4.5	4.6	4.7	4.7	4.7	4.7	4.7	4.7
3	-	4.1	3.7	3.6	4.5	4.6	4.7	4.6	4.7	4.7	4.7	4.7
4	-	4.1	3.6	3.6	4.5	4.5	4.7	4.6	4.6	4.7	4.7	-
5	-	4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.7	4.7	4.7	-
6	-	4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.7	4.7	4.7	-
7	-	4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.7	4.7	4.7	-
8	-	a4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.6	4.7	4.7	a4.8
9	-	a4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.6	4.7	4.7	-
10	-	a4.1	3.6	3.6	4.5	4.6	4.7	4.6	4.6	4.7	4.7	-
11	-	a3.9	3.6	3.6	4.5	4.6	a4.7	4.6	4.6	4.7	4.6	-
12	-	a3.9	3.6	3.6	4.5	4.6	a4.7	4.6	4.6	4.7	4.6	-
13	-	3.9	3.6	3.6	4.5	a4.6	a4.7	4.6	4.6	4.6	4.6	-
14	-	3.8	3.6	3.6	4.5	a4.6	a4.7	4.6	4.6	4.6	4.6	a4.9
15	-	3.8	3.6	3.6	4.5	4.6	a4.7	4.6	4.6	4.6	4.6	-
16	-	3.8	3.6	3.6	4.6	4.6	a4.7	4.6	4.6	4.6	4.6	a5.2
17	-	3.8	3.6	3.6	4.6	4.6	a4.7	4.6	4.6	4.6	4.6	a5.4
18	-	3.8	a3.6	3.6	4.6	4.6	4.7	4.6	4.6	4.6	4.6	-
19	-	3.8	a3.6	3.6	4.6	4.6	4.7	4.6	-	4.6	4.6	-
20	-	3.8	3.6	3.6	4.6	4.6	4.7	4.6	a4.6	4.6	4.6	-
21	-	3.8	3.6	4.1	4.6	4.6	4.7	4.6	-	4.6	4.6	-
22	-	3.8	3.6	4.2	4.6	4.6	4.7	a4.6	-	4.6	4.6	-
23	-	3.7	3.6	4.3	4.6	4.6	4.7	a4.6	a4.7	4.6	4.6	-
24	-	3.7	3.6	4.3	4.6	4.6	4.7	a4.6	-	4.6	4.6	a5.3
25	4.7	3.7	3.6	4.3	4.6	4.6	4.7	a4.6	4.7	4.6	4.6	-

a Estimated, interpolated, or partly estimated.

-63a-
Discharge, in second-feet, of Giffen Springs
At Toyahvale, Tex.--Continued
1931-33

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1931-32	-	-	-	-	-	-	-	-	-	-	-	-
26	4.7	3.7	3.6	4.5	4.6	4.6	4.7	a4.6	4.7	4.6	4.6	} a5.3
27	4.7	3.7	3.6	4.5	4.6	4.6	4.7	a4.6	4.7	4.6	4.6	
28	4.7	3.7	3.6	4.5	4.6	4.6	4.7	a4.6	4.7	4.6	4.6	
29	4.6	3.7	3.6	4.5	4.6	4.6	4.7	4.6	4.7	-	a4.7	
30	4.6	3.7	3.6	4.5	-	4.7	4.7	4.6	4.7	a4.6	a4.7	
31	4.5	-	3.6	4.5	-	4.7	-	4.6	-	-	4.7	

a Estimated, interpolated, or partly estimated.

Discharge, in second-feet, of Giffen Springs at Toyahvale, Tex., 1931-33-Contd.

Day 1932-33	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1					4.9		5.1	4.9	a5.0	4.9	4.9	a5.3
2					4.9		5.1	4.9	a5.0	4.9	4.7	a5.4
3					4.9		5.1	4.9	a5.0	4.9	4.7	-
4					a4.9		5.0	4.9	5.0	4.9	4.7	-
5					a4.9		5.0	4.9	5.0	4.9	4.7	-
6					a4.9		5.0	4.9	5.0	4.9	4.7	-
7					a4.9		5.0	4.9	5.0	4.9	4.7	-
8					a4.9		5.0	4.9	5.0	a4.9	4.7	-
9					a4.9		5.0	4.9	5.0	a4.9	4.7	-
10					a4.9		5.0	4.9	5.0	a4.9	4.7	-
11					4.9		5.0	4.9	5.0	a4.9	4.7	-
12					4.9		5.0	4.9	5.0	a4.9	4.7	-
13					4.9		5.0	4.9	5.0	a4.9	4.7	-
14					4.9		5.0	4.9	4.9	4.9	4.7	-
15					4.9		5.0	4.9	4.9	4.9	4.7	-
16					4.9		5.0	4.9	4.9	4.9	4.6	-
17	6.1	5.9	5.6	5.3			5.0	4.9	4.9	4.9	4.6	-
18							a5.0	4.9	4.9	4.9	4.6	-
19							a5.0	4.9	4.9	4.9	4.6	-
20							a5.0	4.9	4.9	4.9	4.6	-
21							a5.0	5.0	4.9	4.9	4.6	-
22							a5.0	5.0	4.9	4.9	4.6	-
23					5.0		a5.0	5.0	4.9	4.9	4.6	-
24							a5.0	5.0	4.9	4.5	4.6	-
25							a5.0	5.0	4.9	4.5	4.6	-
26							5.1	5.0	4.9	4.3	4.6	-
27							5.1	5.0	4.9	4.3		-
28							5.1	5.0	4.9	4.3		-
29							5.1	4.9	4.9	4.6		-
30							5.1	4.9	4.9	4.9		-
31				4.9			-	5.0	-	4.9		-

a Estimated, interpolated, or partly estimated.

Discharge, in second-feet, of Giffen Springs at Toyahvale, Tex., 1931-33-Contd.

Month	Maximum	Minimum	Mean	Run-off in acre feet
1931-32				
October 25-31	4.7	4.5	4.64	64
November	4.3	3.7	3.89	231
December	3.7	3.6	3.61	222
January	4.5	3.6	3.83	239
February	4.6	4.5	4.55	262
March	4.7	4.6	4.61	283
April	-	-	4.70	280
May	4.7	4.6	4.61	283
June	-	-	4.65	277
July	4.7	4.6	4.64	295
August	4.7	4.6	4.64	285
September	-	-	5.06	301
The period				3,010
1932-33				
October	-	-	6.10	375
November	-	-	5.90	351
December	-	-	5.60	344
January	-	-	5.29	325
February	-	-	4.94	274
March	-	-	5.07	312
April	5.1	4.9	5.00	298
May	5.0	4.9	4.94	304
June	5.0	4.9	4.94	294
July	4.9	4.3	4.81	296
August	-	-	4.70	289
September 1-2	5.4	5.3	5.35	21
The period				3,480

a Estimated, interpolated, or partly estimated.

Discharge, in second-feet, of San Solomon Springs
at Toyahvale, Tex.

1931-33

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1931-32												
1	—	b31			30	b36	34	b32	b32	32	b34	b42
2	—	31			30	36	34	b32	b32	32	b33	b43
3	—	31			30	36	34	b32	32	b32	32	43
4	—	31			30	36	34	b32	32	b33	32	b44
5	—	30	a37		30	35	34	32	32	b33	32	b48
6	—	30			30	35	b34	32	32	b33	32	b49
7	—	30			30	35	b33	32	33	b33	b32	b54
8	—	b30			30	35	33	33	33	33	b32	b58
9	—	b30			30	35	33	33	b34	b33	b32	b60
10	—	b30		b35	30	35	33	33	b34	33	32	62
11	—	b30			30	35	b32	32	b34	33	b32	63
12	—				30	b34	b32	32	b33	33	32	64
13	—				30	b34	b32	32	b33	b33	32	64
14	b32				30	b34	b32	32	b33	b32	32	b65
15	32	a38			30	34	b32	32	b33	b32	32	b65
16	32				30	34	b32	32	b33	b33	32	b64
17	32				30	34	b32	32	b33	b33	32	b64
18	32				30	34	b32	32	33	b33	32	b64
19	32		a36		30	33	b32	32	b33	b33	32	b63
20	32	b38			30	33	b31	32	b32	b33	32	b62
21	32	b38		a32	30	33	b31	32	b32	b33	33	b57
22	b32				30	33	b31	32	b32	b33	33	b56
23	b32	a38		a30	30	33	b31	32	b32	33	b33	56
24	32				31	33	31	32	b32	b33	33	56
25	32				32	b33	31	32	32	b33	b33	55
26	32			a30	34	b33	32	32	b32	b33	33	55
27	32	a37	a35	30	36	33	32	32	b32	b33	33	54
28	32		a35	30	36	b33	32	32	32	b33	32	b54
29	31		b35	30	36	b33	32	32	32	b34	b32	b60
30	b31		a35	30	—	33	32	32	32	34	b37	b66
31	b31	—	a35	30	—	33	—	32	—	b34	40	—

a Estimated

b Partly estimated or interpolated.

Discharge, in sedon-feet, San Solomon Springs, at Toyahvale, Tex., 1931-33-
Continued

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1932-33												
1	66	66	56	b46	41	40	36	b36		b34	b34	48
2	b68	66	55	b46	41	40	36	b36	-	34	35	51
3	b69	66	55	b45	41	40	36	b36	-	a34	35	54
4	b70	65	54	b45	41	40	36	b36	-	.	34	55
5	70	b65	54	b45	40	40	36	b36	.	a36	34	55
6	b70	64	54	b44	b40	40	b36	36			34	55
7	b71	64	53	b44	40	39	36	36	.	a38	34	55
8	b71	64	53	b44	40	39	36	36		38	34	55
9	b71	64	53	b43	40	39	b36	37	a36	38	34	55
10	b70	64	52	b43	40	39	b36	37		38	35	b55
11	b70	64	52	b43	40	38	36	37	-	38	35	56
12	b70	64	52	b43	40	38	36	37		37	35	56
13	b70	64	52	b43	40	38	36	37		37	35	56
14	b70	64	52	b43	40	38	36	37		37	35	56
15	70	64	52	b42	40	38	36	37		37	35	56
16	70	63	51	b42	40	38	36	37		37	35	56
17	70	63	50	b42	40	38	36	37	b36	37	35	56
18	70	62	49	b42	40	38	36	37	35	37	b35	55
19	69	62	48	b42	41	38	36	37	34	37	35	55
20	69	62	47	b42	41	38	36	37	34	37	35	55
21	69	b61	47	b41	41	37	b36	37	34	37	35	55
22	68	60	47	41	41	37	b36	37	b34	a37	35	56
23	68	60	46	41	41	37	36	37	b34	b38	35	56
24	68	59	46	41	41	37	36	36	b35	b38	35	55
25	68	59	b46	41	41	37	b36	36	b35	b38	35	55
26	68	58	b46	41	41	37	36	36	b35	b37	35	b55
27	67	58	b46	41	41	37	36	36	b35	a37	35	
28	67	57	b46	41	40	b37	36	36	b35	a36	35	a54
29	67	57	b46	41	-	b36	36	36	b34	b35	40	
30	66	56	b46	41	-	b36	36	b36	b34	34	41	
31	66	-	b46	41	-	b36	-	b36	-	34	44	

a. Estimated.

b. Partly estimated or interpolated.

Discharge, in second-feet, of San Solomon Springs, at Toyahvale, Tex., 1931-33.
Continued

Day	Oct.	Nov.	Dec.	Day	Oct.	Nov.	Dec.	Day	Oct.	Nov.	Dec.
1933				1933				1933			
1	a50	44	b39	11	a50	41	41	21	46	40	a40
2		43	b39	12		41	41	22	46	40	40
3		43	40	13		41	41	23	46	40	40
4		43	b40	14		41	41	24	46	40	40
5		42	40	15		41	41	25	45	40	39
6		42	40	16	46	41		26	45	40	b39
7		42	41	17	b46	41		27	45	40	39
8		42	41	18	b46	41	a40	28	45	40	39
9		42	41	19	46	40		29	44	39	39
10		42	41	20	46	40		30	44	39	39
								31	44	-	38
Month				Maximum	Minimum	Mean	Run-off in acre-feet				
1931-32											
October 14-31				32	31	31.8	1,140				
November						35.0	2,080				
December						36.2	2,230				
January						33.3	2,050				
February						30.9	1,780				
March					36	30	30.9				
April					36	33	34.1				
May					34	31	32.3				
June					33	32	32.1				
July					34	32	32.5				
August					34	32	32.9				
September					40	32	32.7				
The period				66	42	57.0	3,390				
							24,600				

a. Estimated.

b. Partly estimated or interpolated.

Discharge, in second-feet, of San Solomon Springs at Toyahvale, Tex., 1931-33-
Continued

Month	Maximum	Minimum	Mean	Run-off in acre-feet
1932-33				
October	71	66	68.9	4,240
November	66	56	62.2	3,700
December	56	46	50.1	3,080
January	46	41	43.6	2,620
February	41	40	40.5	2,250
March	40	36	38.1	2,340
April	36	36	36.0	2,140
May	37	36	36.5	2,240
June	-	34	35.3	2,100
July	38	34	36.6	2,250
August	44	34	35.4	2,180
September	-	43	54.8	5,260
The year	71	34	44.7	32,400
1933				
October	-	44	47.4	2,910
November	44	39	41.0	2,440
December	41	38	40.0	2,460
The period				7,810

District measurements of main canal of Reeves County Water Improvement
District No. 1 to determine losses from seepage from source to end, near
Balmorhea, Tex., 1931, 1933

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)				Gain or loss in section	Total gain or loss
				Main stream	Tribu- tary	Diver- sion			
1931									
Oct. 27	Main canal	500 feet below Solomon Spring	0	32.0	-	-	-	-	-
27	Carpenter take-out		.8	-	-	2.81	-	-	-
27	Giffin Spring		1.4	-	1.34	-	-	-	-
27	Reservoir take-out		1.4	-	-	2.83	-	-	-
27	Henry Jones take-out		1.4	-	-	.40	-	-	-
27	North canal take-out		2.2	-	-	.27	-	-	-
27	Gate leakage	Total leakage between points on main canal	-	-	-	.12	-	-	-
27	Main canal	Crenshaw garage, Balmorhea	4.0	25.3	-	-	-1.6	-	-1.6
27	Walker take-out		6.0	-	-	.61	-	-	-
27	Gate leakage	Total leakage between points on main canal	-	-	-	.10	-	-	-
28	Main canal	Highway crossing	7.0	21.4	-	-	-3.2	-	-4.8
28	Highway ditch take-out		7.0	-	-	5.94	-	-	-
28	Sol Mayer take-out		8.0	-	-	5.20	-	-	-
28	Saragosa canal		8.4	-	1.98	-	-	-	-
28	Siphon ditch take-out		8.4	-	-	8.98	-	-	-
28	Gate leakage	Total leakage between points on main canal	-	-	-	.60	-	-	-
28	Main canal	150 feet below Siphon ditch	8.4	1.42	-	.70	-1.2	-	-6.0
28	Gate leakage	Total leakage between points on main canal	-	-	-	-	-	-	-
28	Main canal	$\frac{1}{2}$ mile above end of system	11.4	.38	-	-	-.3	-	-6.3
1933									
Jan. 11	Main canal	500 feet below Giffin canal junction	1.8	48.7	-	-	-	-	-
11	North spill	Knapp's corner	3.1	-	-	26.3	-	-	-
11	Main canal	Balmorhea Hotel	4.7	19.9	-	-	-2.5	-	-2.5
11	Gate leakage	Total of 2 leaks	-	-	-	.1	-	-	-
11	West Sandia canal	Gage	5.6	-	1.8	-	-	-	-
11	Main canal	Bro gajo	6.0	22.2	-	-	4.6	-	-1.9
11	Lateral diversion		6.6	-	-	.8	-	-	-
11	Main canal		7.1	20.2	-	-	-1.2	-	-3.1
11	Gate leakage	Total of 3 leaks	-	-	-	.2	-	-	-
11	Main canal		8.1	21.0	-	-	-2.1	-	-2.1
12	Siphon ditch	Weir 75 feet below main canal	9.6	-	-	1.8	-	-	-

Discharge measurements of main canal of Reeves County Water Improvement
District No. 1 to determine losses from seepage from source to end, near
Balmorhea, Tex., 1931, 1933--Continued

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)				
				Main stream	Tribu- tary	Diver- sion	Gain or loss in section	Total gain or loss
1933								
Jan. 12	Main canal		9.7	17.1	-	-	-2.1	-4.2
12	do.		11.1	17.4	-	-	4 .3	-3.9
Mar. 13	do.	500 feet below Giffin canal junction	1.8	11.3	-	-	-	-
13	do	Wigley road crossing	2.8	11.4	-	-	4 .1	4 0.1
13	Gate leakage	Knepp's corner	3.1	-	-	.1	-	-
13	Main canal	400 feet above Highway Garage	4.7	11.0	-	-	- .3	- .2
13	West Sandia canal	50 feet above highway	5.6	-	1.3	-	-	-
13	Main canal	1 mile below Brogado	6.3	11.9	-	-	- .4	- .6
13	do.	Highway crossing	7.7	11.9	-	-	0	- .6
13	Experiment farm spill	do.	8.3	-	.6	-	-	-
13	Main canal	Above Saragosa canal junction	9.6	12.5	-	-	0	- .6
13	do.	$\frac{1}{2}$ mile below Siphon ditch	10.1	8.3	-	-	-	-
13	do.	Saragosa road crossing	11.1	8.8	-	-	4 .5	4 .5
14	do.	500 feet below Giffin canal junction	1.8	8.2	-	-	-	-
14	Gate leakage		3.0	-	-	.1	-	-
14	do.		3.7	-	-	.1	-	-
14	Main canal	400 feet above Highway Garage	4.7	7.8	-	-	- .2	- .2

Discharge measurements of laterals of Reeves County Water Improvement
District No. 1, to determine losses from seepage near Balmorhea, Tex.,
1931

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)			
				Main stream	Diver- sion	Gain or loss in section	Total gain or loss
Oct. 27	Carpenter take-out	Point of diversion	0	2.81	-	-	-
27	do.	Point of delivery	1.0	2.73	-	-0.08	-0.08
27	Reservoir take-out	Point of diversion	0	2.83	-	-	-
27	do.	Confluence with reservoir creek	1.0	2.33	-	- .50	- .50
27	Highway ditch	Point of diversion	0	9.32	-	-	-
27	Mills ditch	do.	.8	-	5.91	-	-
27	Highway ditch	Below Mills Ditch take-out	.8	3.80	-	/ .39	/ .39
27	do.	Point of delivery to Mayer farm	1.3	3.84	-	/ .04	/ .43
28	Siphon ditch	Point of diversion	0	9.32	-	-	-
28	do.	Point of delivery to Fane Down farm	2.3	6.19	-	-3.13	-3.13
Nov. 16	Moore canal	300 feet below Moore dam	0	4.29	-	-	-
16	do.	Pecos Valley Southern Ry. crossing	.5	4.35	-	/ .06	/ .06
16	Saragosa canal	1,000 feet below diversion dam	0	1.97	-	-	-
16	do.	Weir	.5	1.68	-	- .29	- .29
18	Giffin Spring canal	do.	0	3.80	-	-	-
18	do.	Siphon	.8	4.06	-	/ .26	/ .26

Discharge measurements of reservoir outlet canal of Reeves County Water
Improvement District No. 1 to determine losses from seepage from release gate
at reservoir to junction with main canal, near Balmorhea, Tex., 1932-33

Date	Stream and diversion	Location	Distance from initial point (miles)	Discharge (second-feet)			
				Main stream	Diversion	Gain or loss in section	Total gain or loss
1932							
July 22	Outlet canal	0.1 mile below release gate	0.1	52.2	-	-	-
22	Gate leakage	0.3 mile below release gate	.3	-	0.4	-	-
22	Outlet canal	3.5 miles below release gate	3.5	48.0	-	-3.8	-3.8
26	do.	0.1 mile below release gate	.1	42.1	-	-	-
26	Gate leakage	0.3 mile below release gate	.3	-	.3	-	-
26	Outlet canal	0.7 mile below release gate	.7	40.9	-	- .9	- .9
26	do.	1.7 miles below release gate	1.7	41.1	-	/ .2	- .7
26	do.	2.4 miles below release gate	2.4	39.8	-	-1.3	-2.0
26	do.	50 feet above main canal	3.5	41.5	-	/1.7	- .3
Aug. 17	do.	0.1 mile below release gate	.1	1.8	-	-	-
17	do.	0.4 mile below release gate	.4	2.1	-	/ .3	/ .3
17	do.	2.6 miles below release gate	2.6	1.6	-	- .5	- .2
1933							
July 20	do.	0.2 mile below release gate	.2	14.2	-	-	-
20	do.	0.4 mile above main canal	3.1	11.6	-	-2.6	-2.6
26	do.	0.2 mile below release gate	.2	7.2	-	-	-
26	do.	0.6 mile above main canal	2.9	6.0	-	-1.2	-1.2

Discharge, in second-feet, of West
Sandia Spring at Balmorhea, Tex.

1931-32

Day	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.	Sept.
1931-32												
1			al.1	1.1		1.3	1.5	1.2		1.1	al.1	a b.1.1
2			al.1	1.1		1.5	1.2	1.1		1.1	al.1	a b.1.3
3			1.2	1.1	al.1	1.3	1.2	1.1		1.1	al.1	b.1.4
4			1.2	1.1		1.4	1.2	1.1	al.1	1.1	1.1	a b.1.4
5			1.2	1.1		1.3	1.2	1.0		1.1	1.1	a b.1.5
6			1.2	1.1	al.1	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.5
7			1.2	1.1	al.1	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.6
8			1.2	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
9			1.2	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
10			1.2	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
11			1.2	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
12			1.2	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
13			1.1	1.1	al.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
14			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
15			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
16			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
17			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
18			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
19			1.1	1.1	1.2	1.3	1.2	1.0	al.1	1.1	1.1	a b.1.7
20			1.1	1.1	al.2	1.3	al.2		al.1	1.1	1.0	1.7
21			1.1	1.1	al.3	1.3	al.2		al.1	1.1	1.0	1.7
22			1.1	1.1	al.3	1.3	al.3		al.1	1.1	1.0	1.7
23			1.1	1.1	al.3	1.3	al.3		al.1	1.1	1.0	1.7
24			al.1	1.1	al.3	1.2	b1.3		al.1	1.0	1.0	a.1.7
25			al.1	1.1	al.3	1.2	b1.3	al.1	1.2	1.0	1.0	a.1.6
26			al.1	1.1	al.3	1.2	1.2	1.2	1.2	1.0	1.0	a.1.6
27			al.1	1.1	al.3	1.2	1.2	1.2	1.2	1.0	1.0	a.1.6
28			al.1	1.1	al.3	1.2	1.2	1.2	1.2	1.0	1.0	a.1.6
29			al.1	1.1	1.3	al.3	1.2	1.2	1.2	1.0	1.0	a.1.6
30			al.1	1.1	1.3	al.3	1.2	1.2	1.2	1.0	1.0	a.1.6
31			1.1	al.1	-	al.3	-		-	al.0	al.0	a.1.7

a Discharge estimated or partly estimated.

b May include some storm run-off or irrigation waste water.

Discharge, in second-feet, of West Sandia Spring at Balmorhea, Tex., 1932-35-Con.

Day 1932-33	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	Apr.	May	June	July	Aug.
1	a 1.3	1.8	1.7		b 1.7	b 1.6	b 1.3	b 1.2		1.1	1.1
2		1.8	1.7		b 1.7	b 1.6	b 1.3	b 1.2	a 1.2	1.1	1.1
3		1.8	1.6		b 1.7	b 1.6	b 1.3	b 1.2		1.1	1.1
4		1.8	1.6		b 1.7	b 1.6	b 1.3	b 1.2		1.1	1.1
5		1.8	1.6	a 1.7	b 1.7	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
6		1.8	1.6		b 1.7	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
7		1.7	1.6		b 1.7	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
8		1.7	1.7		b 1.7	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
9		1.7	1.7		b 1.7	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
10	a 1.9	1.7	1.7	b 1.7	b 1.6	b 1.6	b 1.3	b 1.2	b 1.2	1.1	1.1
11		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
12		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
13		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
14		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
15		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
16		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
17	b 1.9	1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
18		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
19		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
20		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
21		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
22		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
23	a 1.9	b 1.7	b 1.7	1.5	b 1.6	b 1.4	b 1.3	b 1.2	b 1.2	1.1	1.1
24		b 1.7	b 1.7	1.5	b 1.6	b 1.4	b 1.3	b 1.2	b 1.2	1.1	1.1
25		b 1.7	b 1.7	1.5	b 1.6	b 1.4	b 1.3	b 1.2	b 1.2	1.1	1.1
26		b 1.7	b 1.7	1.5	b 1.6	b 1.4	b 1.3	b 1.2	b 1.2	1.1	1.1
27		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
28		1.7	1.7	1.7	b 1.6	b 1.5	b 1.3	b 1.2	b 1.2	1.1	1.1
29	1.9	1.7	a 1.7							1.1	1.1
30	1.8	1.7		b 1.8	-	b 1.5	1.2		b 1.1	1.1	-
31	1.8	-		b 1.7	-	b 1.3	-		b 1.1	1.1	-

a Estimated
b Partly estimated.

Discharge, in second-feet, of West Sandia Spring at Balmorhea, Tex., 1932-33-Con.

Month	Maximum	Minimum	Mean	Run-off in acre feet
1931-32				
November 17-30	1.1	1.1	1.10	31
December	1.2	1.1	1.13	69
January	1.1	1.1	1.10	68
February	1.4	1.1	1.21	70
March	1.4	1.2	1.29	79
April	1.3	1.2	1.21	72
May			1.11	68
June			1.12	67
July	1.1	1.0	1.06	65
August	1.1	1.0	1.03	63
September	1.7	1.1	1.61	96
The period				748
1932-33				
October			1.89	116
November	1.8	1.7	1.72	102
December			1.68	103
January			1.66	102
February	1.7	1.6	1.63	91
March	1.6	1.3	1.47	90
April	1.3	1.2	1.28	76
May			1.20	74
June			1.15	68
July	1.1	1.1	1.10	68
August 1-23	1.1	1.0	1.06	59
The period				949

a Estimated.

b Partly estimated.

Discharge measurements of West Sandia Creek to determine seepage from 4,000 feet above to gage 0.8 mile east of Balmorhea, Tex., Oct. 17, 1932

Stream	Location	Distance above gage (feet)	Discharge (second-feet)			
			Main stream	Tributary	Gain or loss in section	Total gain or loss
West Sandia Creek	500 feet above West Sandia spring	4,000	0.2	-	-	-
do.	80 feet below West Sandia spring	3,620	.6	-	0.4	0.4
Canal wasteway	500 feet below West Sandia spring	3,200	-	0.1	-	-
West Sandia Creek	Gage	0	2.2	-	1.5	1.9

Discharge, in second-feet, of East Sandia Spring at Balmorhea, Tex.
1931-32

Month	Maximum	Minimum	Mean	Run-off in acre feet
November 23-30 1931-32	1.1	1.1	1.10	17
December	1.1	1.0	1.06	65
January	1.1	1.1	1.10	68
February	-	1.1	1.22	70
March	-	-	1.27	73
April	-	-	1.20	71
May	-	-	1.20	74
June	1.2	.9	1.05	62
July	-	-	1.07	66
August	-	1.0	1.04	64
September	-	-	1.35	30
The period				715
1932-33				
October	-	-	1.35	83
November	1.3	1.2	1.23	73
December	-	-	1.21	74
January	-	-	1.23	79
February	1.4	1.2	1.34	74
March	-	-	1.32	81
April	-	-	1.30	77
May	-	-	1.16	71
June	-	-	1.09	65
July	1.0	1.0	1.00	61
August	1.1	.9	.92	57
September 1-23	-	-	1.10	50
The period				845

Note. - Gage is 10 miles above United States Highway 290 crossing.

Discharge measurements of Cherry Canyon to determine losses
from seepage from a point 1.5 miles above to a point 2.5
miles below gage near Toyahvale, Tex., 1932

Date	Location	Distance from initial point (miles)	Discharge (second-foot)		
			Main stream	Gain or loss in section	Total gain or loss
Sept. 15	Gage	0	15.5	-	-
15	500 feet above Kingston line fence	2.0	0	-13.5	-13.5
21	1.5 miles above gage	-1.5	1.2	-	-
21	Gage	0	5.0	+ 3.8	+ 3.8
Oct. 7	Gage	0	31.2	-	-
7	2.5 miles below gage	2.5	0	-31.2	-31.2

Note. - Gage is 10 miles above United States Highway 290 crossing.

Discharge measurements of Limpia Creek to determine losses from
seepage from a point 12.3 miles above to a point 40.2 miles below
Fort Davis, Tex., 1932-33

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)				
				Main stream	Tribu- tary	Diver- sion	Gain or loss in section	Total gain or loss
1932								
Oct. 8	Limpia Creek	0.8 mile below Wild Rose Canyon	14.7	31.0	-	-	-	-
8	Short Canyon	100 feet above mouth	15.8	-	1.8	-	-	-
8	Limpia Creek	500 feet below old Limpia post office	18.2	46.0	-	-	13.2	13.2
8	do.	3 miles below old Limpia post office	21.2	41.8	-	-	- 4.2	9.0
8	Horse Thief Canyon	Mouth	27.7	-	1.2	-	-	-
8	Runey Canyon	do.	27.9	-	1.5	-	-	-
8	Limpia Creek	Jeff ranch house	28.2	65.1	-	-	20.6	29.6
8	do.	9 miles below Jeff ranch	37.2	4.2	-	-	-60.9	-31.3
8	do.	12 miles below Jeff Ranch	40.2	0	-	-	- 4.2	-35.5
18	do.	12.3 miles above old Fort Davis lane	-12.3	.1	-	-	-	-
18	do.	12 miles above old Fort Davis lane	-12.0	0	-	-	- .1	- .1
18	do.	11.3 miles above old Fort Davis lane	-11.3	.2	-	-	.2	.1
18	do.	11 miles above old Fort Davis lane	-11.0	0	-	-	- .2	- .1
18	do.	10.3 miles above old Fort Davis lane	-10.3	.5	-	-	.5	.4
18	do.	9.8 miles above old Fort Davis lane	- 9.8	0	-	-	- .5	- .1
18	do.	9.7 miles above old Fort Davis lane	- 9.7	.1	-	-	.1	0
18	do.	7.9 miles above old Fort Davis lane	- 7.9	.6	-	-	.5	.5
18	do.	6 miles above old Fort Davis lane	- 6.0	5.7	-	-	5.1	5.6
18	Side canyon	5.9 miles above old Fort Davis lane	- 5.9	-	.5	-	-	-
18	Limpia Creek	3.7 miles above old Fort Davis lane	- 3.7	5.7	-	-	- .5	5.1
18	do.	1.3 miles above old Fort Davis lane	- 1.3	5.3	-	-	- .4	- 4.7
18	Grayson diversion	Old Fort Davis lane	0	-	-	0.5	-	-
18	Limpia Creek	First Fort Davis-Toyahvale crossing	1.2	3.8	-	-	- 1.0	3.7
19	do.	do.	1.2	3.8	-	-	7.6	-
19	Side canyon	2.1 miles below old Fort Davis lane	2.1	-	.4	-	7.2	-
19	do.	do.	2.1	-	.2	-	-	-
19	Limpia Creek	4.2 miles below old Fort Davis lane	4.2	8.0	-	-	3.6	7.3
19	Side canyon	4.8 miles below old Fort Davis lane	4.8	-	.5	-	-	-
19	Limpia Creek	7.2 miles below old Fort Davis lane	7.2	8.8	-	-	.3	7.6
19	do.	10.2 miles below old Fort Davis lane	10.2	8.4	-	-	- .4	- 7.2
19	Frazier Canyon	Mouth	10.3	-	.4	-	-	-
19	Limpia Creek	Upper end of Wild Rose Canyon	11.7	9.4	-	-	.6	7.8
19	do.	Lower end of Wild Rose Canyon	14.0	12.9	-	-	3.5	11.3
19	Short Canyon	Mouth	15.8	-	.1	-	-	-

Discharge measurements of Limpia Creek to determine losses from
seepage from a point 12.3 miles above to a point 40.2 miles below
Fort Davis, Tex., 1932-33--Continued

Date	Stream or diversion	Location	Distance from initial point (miles)	Discharge (second-feet)			Gain or loss	
				Main stream	Tribu- tary	Diver- sion	loss in section	gain or loss
1932								
Oct. 19	Limpia Creek	Old Limpia post office	18.2	14.0	-	-	/ 1.0	/12.3
Nov. 1	do.	do.	18.2	8.2	-	-	-	-
1	do.	3 miles below old Limpia post office	21.2	5.4	-	-	- 2.8	- 2.8
1	do.	5.9 miles below old Limpia post office	24.1	4.6	-	-	- .8	- 3.6
1	do.	Jeff ranch house	28.2	6.2	-	-	/ 1.6	- 2.0
1	do.	1 mile below Jeff ranch house	29.2	0	-	-	- 6.2	- 8.2
21	do.	Upper end of Wild Rose Canyon	11.7	4.0	-	-	-	-
21	do.	Lower end of Wild Rose Canyon	14.0	6.1	-	-	/ 2.1	/ 2.1
21	do.	Old Limpia post office	18.2	4.7	-	-	- 1.4	/ .7
21	do.	3 miles below old Limpia post office	21.2	2.0	-	-	- 2.7	- 2.0
21	do.	6 miles below old Limpia post office	24.2	1.1	-	-	- .9	- 2.9
21	do.	Jeff ranch house	28.2	2.4	-	-	/ 1.3	- 1.6
21	do.	0.5 mile below Jeff ranch house	28.7	0	-	-	- 2.4	- 4.0
1933								
Aug. 3	do.	75 feet below mouth of Short Canyon	15.8	12.6	-	-	-	-
3	do.	Old Limpia post office	18.2	4.0	-	-	- 3.6	- 3.6
3	do.	0.5 mile below old Limpia post office	18.7	0	-	-	- 4.0	-12.6

Miscellaneous discharge measurements near
Balmorhea, Tex., during the years ending
Sept. 30, 1932 and 1933

1931				
Oct. 26	Madera Canyon	Toyah Creek	13.3 miles above Toyahvale, Tex.	0
1932				
July 18	do.	do.	do.	0
Sept. 2	Toyah Creek	Pecos River	U.S. Highway 290 crossing near Toyahvale, Tex.	463
2	do.	do.	do.	551
7	do.	do.	Old county steel bridge 3 miles below Balmorhea, Tex.	c26,100
29	do.	do.	do.	11,400
1933				
Aug. 8	do.	do.	Flow of springs in creek 2.5 miles east of Hoban station and 14 miles below Saragosa, Tex.	a1.5
July 10	Little Aguja Canyon	Toyah Creek	18 miles above Toyahvale, Tex.	a.05
1931				
Oct. 26	do.	do.	Odell ranch 15 $\frac{1}{2}$ miles above Toyahvale, Tex.	0
1932				
July 18	do.	do.	do.	0
29	do.	do.	do.	0
Aug. 16	do.	do.	do.	0
17	do.	do.	do.	1.32
29	do.	do.	do.	c2,640
Dec. 16	do.	do.	do.	a.06
1933				
Jan. 26	do.	do.	do.	a.05
Mar. 9	do.	do.	do.	0
Apr. 10	do.	do.	do.	0
July 10	do.	do.	do.	0
May 18	South Fork of Little Aguja Canyon	Little Aguja Canyon	1 $\frac{1}{2}$ miles above mouth, near Toyahvale, Tex.	.47
July 10	do.	do.	do.	.24
1932				
Sept.--	do.	do.	1 $\frac{1}{2}$ miles above mouth, near Toyahvale, Tex.	1,410
1931				
Oct. 26	do.	do.	$\frac{1}{2}$ mile above mouth, near Toyahvale, Tex.	a.2
1932				
Aug. 16	do.	do.	do.	a.30
17	do.	do.	do.	a.32
Sept. 1	do.	do.	do.	6.45
Dec. 16	do.	do.	do.	a.33
1933				
Jan. 26	do.	do.	do.	a.16
Mar. 9	do.	do.	do.	a.13
Apr. 10	do.	do.	do.	a.20
May 18	do.	do.	do.	a.15
July 10	do.	do.	do.	0
Mar. 9	Big Aguja Canyon	Toyah Creek	200 feet below spring, 3 miles above Texas & Pacific Ry. Co. dam, 15.6 miles above Toyahvale, Tex.	1.02
9	do.	do.	500 feet above Texas & Pacific Ry. Co. reservoir, 15.6 miles above Toyahvale, Tex.	1.69

a Estimated. c Determined by slope-area method.

Miscellaneous discharge measurements near
Balmorhea, Texas during the years ending
Sept. 30, 1932 and 1933-Continued

Date	Stream	Tributary to or diverting from	Locality	Discharge
1933				Sec. -ft.
May 18	Big Aguja Canyon	Toyah Creek	On spillway of Texas & Pacific Ry. Co. dam 12.6 miles above Toyah- vale, Tex.	a.14
July 10	do	do	do	a.25
1931				
Oct. 26	do	do	3/4 mile below Texas & Pacific Ry. Co. dam 11.8 miles above Toyahvale, Texas	ad.1
1932				
July 18	do	do	do	ad.32
Aug. 16	do	do	do	ad.3
Sept. 1	do	do	do	f18.7
3	do	do	do	d45.4
6	do	do	do	d151
7	do	do	do	d4,360
13	do	do	do	d8.6
20	do	do	do	d2.56
Oct. 6	do	do	do	d10.3
Dec. 16	do	do	do	d1.13
1933				
Jan. 26	do	do	do	d1.36
Mar. 9	do	do	do	d.12
Apr. 10	do	do	do	ad.15
May 18	do	do	do	ad.09
July 10	do	do	do	d.17
May 18	Duncan Diversion	Big Aguja Canyon	Just below point of diversion, 11.8 miles above Toyahvale, Texas	1.05
July 10	do	do	do	.95
1931				
Oct. 20	Seven Springs	do	State highway 17 about 7 $\frac{1}{2}$ -8 $\frac{1}{2}$ miles south of Toyahvale, Texas	a.10
1932				
July 13	do	do	do	a.13
Aug. 16	do	do	do	a.15
Dec. 16	do	do	do	a.09
1933				
Jan. 26	do	do	do	a.16
Mar. 9	do	do	do	a.04

a Estimated.

d Does not include J. C. Duncan diversion 25 feet below section.

e Lake level normal.

f Lake level 1 foot above normal.

g Flow in low ditch.

h Flow in high ditch.

Miscellaneous discharge measurements near Balmorhea, Texas during
the years ending Sept. 30, 1932 and 1933--

Continued

Date	Stream	Tributary to or diverting from	Locality	Discharge
1933				Sec.-ft
Apr. 10	Seven Springs	Big Aguja Canyon	State highway 17 about 7½-8½ miles south of Toyahvale, Texas	a.36
May 18	do	do	do	a.09
July 10	do	do	do	a.14
1932				
Aug. 1	Madera diversion canal	Toyah Creek	Bridge on State high- way 17, 1¼ mile south of Toyahvale, Tex.	a.34
17	do	do	do	8.19
30	do	do	do	1,930
30	do	do	do	1,650
Sept. 1	do	do	do	3.22
2	do	do	do	81.2
1931				
Oct. 16	Phantom Lake Spring	do	Source, near Toyah- vale, Texas	e12.6
17	do	do	do	f10.3
19	do	do	do	f11.3
21	do	do	do	f10.9
21	do	do	do	f10.0
22	do	do	do	f11.0
22	do	do	do	f10.9
22	do	do	do	f11.1
22	do	do	do	f10.9
22	do	do	do	f11.3
19	Giffin Springs	do	Source, Toyahvale, Texas	g 4.26
20	do	do	do	h 3.82
21	do	do	do	h 3.64
24	do	do	do	h 4.07
24	do	do	do	h 4.03
24	do	do	do	h 4.15

a Estimated.

d Does not include J. C. Duncan diversion 25 feet below section.

e Lake level normal.

f Lake level 1 foot above normal.

g Flow in low ditch.

h Flow in high ditch.

Miscellaneous discharge measurements near
Balmorhea, Tex., during the years ending
Sept. 30, 1932 and 1933--Continued

Date	Stream	Tributary to or diverting from	Locality	Discharge
1931				sec. ft.
Nov. 10	San Solomon Springs	Toyah Creek	Source, Toyahvale, Tex	h30.9
20	do.	do.	do.	g37.7
21	do.	do.	do.	g37.6
21	do.	do.	do.	g38.0
Dec. 29	do.	do.	do.	g35.4
1932				
Jan. 28	do.	do.	do.	h31.2
1933				
July 13	do.	do.	do.	hm28.6
24	do.	do.	do.	h37.6
26	do.	do.	do.	h36.9
1932				
Nov. 6	Saragosa Springs	do.	$\frac{1}{2}$ mile west of Balmorhea, Tex.	9.15
1933				
Jan. 23	do.	do.	do.	8.22
Mar. 14	do.	do.	do.	6.69
May 16	do.	do.	do.	6.36
July 11	do.	do.	do.	5.59
29	Stock streams	do.	Released for stock water from Reeves County Water Improve- ment district No. 1 canal system near Balmorhea, Tex.	.89
29	Sandia Creek	do.	$\frac{3}{4}$ mile east of State Experiment Farm, near Balmorhea, Tex.	a.45
1932				
Sept. 8	Reservoir north spillway	Sandia Creek	2 miles southeast of Balmorhea, Tex.	225
17	do.	do.	do.	.72
Oct. 1	do.	do.	do.	1,450
Aug. 2	Texas & Pacific farm well no. 3	Toyah Creek	Hoban station, 16 miles south of Pecos, Tex.	2.26
2	Texas & Pacific farm well no. 1	do.	do.	1.86
8	Texas & Pacific farm well no. 4	do.	do.	1.95
July 18	Cherry Canyon	do.	5 miles above Jeff Davis-Reeves County line, near Toyahvale, Tex.	0
Aug. 27	do.	do.	do.	3.10
27	do.	do.	do.	1.37
Sept. 15	do.	do.	do.	13.5
21	do.	do.	do.	4.98
27	do.	do.	do.	5.66
29	do.	do.	do.	c5,320
Oct. 7	do.	do.	do.	31.2
Dec. 15	do.	do.	do.	a.10
1933				
Jan. 25	do.	do.	do.	a.13
Mar. 12	do.	do.	do.	a.08
Apr. 10	do.	do.	do.	0
May 18	do.	do.	do.	0
July 7	do.	do.	do.	0

Miscellaneous discharge measurements near
Balmorhea, Tex., during the years ending
Sept. 30, 1932 and 1933--Continued

Date	Stream	Tributary to or diverting from	Locality	Discharge
1933				
July 29	J. L. Moore stock stream	Cherry Canyon	$\frac{3}{4}$ mile above Balmorhea-Toyah road crossing, 6 miles northeast of Balmorhea, Tex.	a.16
1932				
Aug. 5	Youngblood south well	Toyah Creek	75 feet east of Pecos Valley Southern Ry. $13\frac{1}{2}$ miles south of Pecos, Tex.	3.99
	5 Youngblood north well	do.	$\frac{1}{2}$ mile east of Pecos Valley Southern Ry. 13 miles south of Pecos, Tex.	3.04
	5 J. H. Sudbrock well	do.	$\frac{1}{2}$ mile west of Pecos Valley Southern Ry. $13\frac{1}{2}$ miles south of Pecos, Tex.	2.07
	5 John Wendt well	do.	150 feet east of Pecos Valley Southern Ry. $11\frac{1}{2}$ miles south of Pecos, Tex.	2.77
	11 Balmorhea Live- stock Co. well	do.	1 mile east of Pecos Valley Southern Ry., 8 miles south of Pecos, Tex.	.65
	8 W. A. Gardner well	do.	$\frac{3}{4}$ mile west of Pecos Valley Southern Ry., $4\frac{1}{2}$ miles south of Pecos, Tex.	.95
1933				
Aug. 2	Limpia Creek	Barrilla Creek	Lower end of Wild Rose Canyon 14 miles below Fort Davis, Tex.	a.3

a Estimated.

c Determined.

g Flow in low ditch.

h Flow in high ditch.

m Not total flow of springs.

Miscellaneous discharge measurements near
Balmorhea, Tex., during the years ending
Sept. 30, 1932 and 1933--Continued


Date	Stream	Tributary to or diverting from	Locality	Discharge Sec. ft.
1932				
Sept. 26	Limpia Creek	Barrilla Creek	Old Limpia post office, 12 miles south of Toyahvale, Tex.	121
Aug. 13	Southern Crude Oil Co. flowing well	do.	About 25 miles southeast of Pecos, Tex.	11.96
Oct. 28	Humble Thompson flowing well	Leon Creek	9 miles west of Fort Stockton, Tex.	1.35
Aug. 24	Leon Springs and flowing wells	do.	Just above Leon Springs reser- voir, 7 miles west of Fort Stockton, Tex.	16.0
1933				
Aug. 18	do.	do.	do.	18.0
1932				
Aug. 3	Comanche Springs	Comanche Creek	Main canal $\frac{1}{4}$ mile below diversion dam, at Fort Stockton, Tex.	40.9
24	do.	do.	do.	k45.0
Sept. 14	do.	do.	do.	44.3
Oct. 10	do.	do.	do.	46.7
26	do.	do.	do.	47.7
Nov. 10	do.	do.	do.	48.6
22	do.	do.	do.	50.0
Dec. 14	do.	do.	do.	49.1
1933				
Mar. 17	do.	do.	do.	k48.2
Apr. 11	do.	do.	do.	47.8
May 16	do.	do.	do.	47.0
July 2	do.	do.	do.	48.0
19	do.	do.	do.	<u>1/47.6</u>
27	do.	do.	do.	46.8
Aug. 4	do.	do.	do.	45.9
10	do.	do.	do.	44.3
14	do.	do.	do.	<u>1/46.0</u>
18	do.	do.	do.	<u>1/45.1</u>
Sept. 29	do.	do.	do.	<u>1/43.4</u>
1932				
Oct. 28	Adobe Springs	do.	11 miles east of Fort Stockton, Tex.	.37
28	Miracle and two other flowing wells	do.	12 miles northeast of Fort Stockton, Tex.	1.59
28	Tourney no. 1 flow- ing well	do.	do.	5.23
28	Trans-Pecos no. 1 flowing well	do.	12 $\frac{1}{2}$ miles northeast of Fort Stockton, Tex.	.97
26	Six Shooter Draw	Pecos River	2 miles below Tunis Spring, 23 miles east of Fort Stockton, Tex.	2.34
26	Tunis Spring	Six Shooter Draw	21 miles east of Fort Stockton, Tex.	1.73
Nov. 10	do.	do.	do.	1.98

Miscellaneous discharge measurements near
Balmorhea, Tex., during the years ending
Sept. 30, 1932 and 1933-Continued

Date	Stream	Tributary to or diverting from	Locality	Discharge Sec. ft.
1933				
Jan. 13	Tunis Spring	Six Shooter Draw	21 miles east of Fort Stockton, Tex.	1.71
Mar. 17	do.	do.	do.	1.94
May 13	do.	do.	do.	1.73
1932				
Oct. 11	Rio La Luz	Tularosa Valley Basin	Sec. 28, T. 15 S., R. 10 E., at head of Alamogordo community ditch, 1 mile east of La Luz, N. Mex.	10.7
Nov. 6	do.	do.	do.	10.9
Dec. 14	do.	do.	do.	13.9
1933				
Jan. 13	do.	do.	do.	13.2
Feb. 5	do.	do.	do.	12.4
Mar. 7	do.	do.	do.	11.9
Apr. 15	do.	do.	do.	6.60
May 24	do.	do.	do.	6.64
June 10	do.	do.	do.	2.62
July 17	do.	do.	do.	36.1
Aug. 11	do.	do.	do.	8.21
12	do.	do.	do.	6.54
31	do.	do.	do.	3.54
Sept. 20	do.	do.	do.	14.1

- i Well opened to full capacity 20 minutes before starting measurement.
- j Well partly shut off.
- k Spring openings and channels being cleaned.
- l Spring water backed up over spring openings by growth in creek channel.

■ SAN MARTINE




EXPLANATION



○ SPRING

▨ PRINCIPAL KNOWN AREAS OF GROUND WATER INTAKE



MAP OF PART OF TOYAH BASIN, TEXAS
 SHOWING SPRINGS AND AREAS OF GROUND WATER INTAKE
 NEAR BALMORHEA.




By W. N. White

MAP OF THE VICINITY OF BALMORHEA, TEXAS

SHOWING WELLS AND SPRINGS AND APPROXIMATE
SHAPE AND SLOPE OF WATER TABLE

By S. S. Nye

- EXPLANATION -

-  RECORD WELL
-  SPRING
-  LINES OF EQUAL ALTITUDE
OF WATER LEVELS IN WELLS
IN 1930-31.
(ASSUMED DATUM)








FIGURE 2



EXPLANATION



SECTION ALONG LINE A-A'



GEOLOGIC MAP AND SECTION OF BALMORHEA AREA, TEXAS, SHOWING THE RELATIONS BETWEEN THE LARGE SPRINGS AND GEOLOGIC STRUCTURE.

THE LARGE SPRINGS OCCUR AS A RESULT OF FAULTING WHICH HAS BROUGHT DOWN RELATIVELY IMPERMEABLE UPPER CRETACEOUS ROCKS AGAINST CAVERNOUS LOWER CRETACEOUS LIMESTONE